arXiv:1207.5011v1 [math.DG] 20 Jul 2012

CONICAL KÄHLER-EINSTEIN METRIC REVISITED

CHI LI, SONG SUN

Abstract. The purpose of this paper is of several folds 1. Review the recent development of Kähler-Einstein metric with cone singularities along a divisor. 2. Construct Kähler-Einstein metrics using branch covers. 3. Further extend some standard (both analytic and algebraic sides of) theory in the smooth Kähler-Einstein case to the conical setting. In particular, we use Berndtsson's convexity result to reprove Chen's theorem on the lower boundedness of K-energy in the Kähler-Einstein case and generalize it to the singular and conical setting. 4. Propose a degeneration-interpolation method to construct conical Kähler-Einstein metrics. 5. As a motivational example, we prove that there exists a Kähler-Einstein metric on $\mathbb{P}^2$ with cone singularity along a smooth conic (degree 2) curve if and only if the angle is between $\pi/2$ and 2π. When the angle is $2\pi/3$ this provides a Sasaki-Einstein metric on the link of a three dimensional A_2 singularity. 6. Prove a version of Donaldson's conjecture in the toric case using Song-Wang's recent existence result of toric invariant conical Kähler-Einstein metrics.

Contents

1. Introduction 2
2. preliminaries 5
2.1. Space of admissible potentials for conical Kähler metrics 5
2.2. Functional from smooth Kähler-Einstein 7
2.3. Degenerate Complex Monge-Ampère equation on Kähler manifold with boundary 9
3. Obstruction to existence: log-K-stability 11
3.1. log-Futaki invariant and log-K-(semi)stability 11
3.2. log-Mabuchi-energy and log-Futaki-invariant 12
3.3. log-slope stability and log-Fano manifold 15
4. Existence of conical Kähler-Einstein metric 18
4.1. log-Mabuchi-energy and log-Ding-energy 18
4.2. Existence of conical Kähler-Einstein metric 20
4.3. Alpha-invariant and small cone angles 21
4.4. Construct smooth Kähler-Einstein metrics on branch covers 23
5. Special degeneration to Kähler-Einstein varieties 24
5.1. Kähler current on singular varieties 24
5.2. Proof of Theorem 1.4 25
6. A case study: D=degree 2 curve on $X = \mathbb{P}^2$ 27
6.1. Proof of Theorem 1.7 27
6.2. Calabi-Yau cone metrics on three dimensional A_2 singularity 29
7. Convergence of conical KE to smooth KE 30
8. Relations to Song-Wang's work 36
References 40

Date: July 23, 2012.

1. INTRODUCTION

The existence of smooth Kähler-Einstein metrics on a Kähler manifold X is a main problem in Kähler geometry. The problem is of trichotomic nature. The case when the first Chern class of X is negative was solved by Aubin [2] and Yau [65]. The case when the first Chern class is zero follows from Yau's celebrated solution of Calabi conjecture in [65]. However, when the first Chern class is positive, in other words, when the manifold is Fano, there are obstructions for the existence. The study of such obstructions is the main goal of Yau-Tian-Donaldson program.

There are variations or generalizations of problem of smooth Kähler-Einstein metrics. One variation is to consider Kähler-Einstein metrics with cone singularities. This problem was classically studied on the Riemann surfaces ([40, 63, 38]) and was first considered in higher dimension by Tian in [57]. Recently, there are reviving works on the Kähler-Einstein metrics with cone singularities. ([20, 6, 31, 35]) These are partly motivated by Donaldson's project (see [20]) of using conical Kähler-Einstein metric as a continuity method to attack the existence problem of smooth Kähler-Einstein metric on Fano manifolds.

From now on in this paper, we assume X is a smooth Fano manifold. Assume D is a smooth divisor such that $D \sim_{\mathbb{Q}} -\lambda K_X$ for some $\lambda > 0 \in \mathbb{Q}$. Notice that $K_X + D \sim_{\mathbb{Q}} (1-\lambda)K_X$. So (X, D) is log canonical-polarized (resp. log Calabi-Yau, resp. log $\mathbb{Q}$-Fano) when $\lambda > 1$ (resp. $\lambda = 1$, resp. $\lambda < 1$). We will consider the existence problem of singular Kähler-Einstein metric in the class $c_1(X) > 0$ with cone singularity along D. In other words, if the cone angle is $2\pi\beta$, we will be interested in solving the following equation:

$$Ric(\omega) = r(\beta)\omega + (1-\beta)\{D\}. \tag{$*$}$$

By taking cohomological class on both sides, we get the formula

$$r(\beta) = 1 - (1-\beta)\lambda. \tag{1}$$

We will use this notation throughout the paper.

One of our motivation for this paper is to construct smooth Kähler-Einstein metrics using branch covers(see [1], [25] for such kind of constructions). If $D \sim mD_1$ with D_1 being an integral divisor, we can construct branch cover of X with branch locus D.

$$\begin{tikzcd} B \arrow[r, phantom, "\subset"] \arrow[d] & Y \arrow[d, "\pi"] \\ D \arrow[r, phantom, "\subset"] & X \end{tikzcd}$$

Assume we have already constructed an orbifold Kähler-Einstein metric $\widehat{\omega}_{KE}$ on $(X, (1-1/m)D$. Then $\pi^*\widehat{\omega}_{KE}$ is a smooth Kähler-Einstein metric on Y. Note that orbifold Kähler metric can be seen as a special case of conical Kähler metric, i.e. when the cone angle is equal to $2\pi/m$ for some $m \in \mathbb{Z}$. So existence of conical Kähler-Einstein metrics with angle $2\pi/m$ will give rise to smooth Kähler-Einstein metrics.

By the work of [31] and [6], we already have the following existence result. See section 4.2 for discussions of these results.

Theorem 1.1. *([31],[6])*

1. *(log canonical-polarized) If $\lambda > 1$, then there exists a conical Kähler-Einstein metric ω_β on $(X, (1-\beta)D)$ when $0 < \beta < 1 - \lambda^{-1} + \epsilon$ for $0 < \epsilon \ll 1$.*
2. *(log Calabi-Yau) If $\lambda = 1$, then there exists a conical Kähler-Einstein metric ω_β on $((X, (1-\beta)D), c_1(X))$ when $0 < \beta < \epsilon$. Such Kähler-Einstein metric ω_β has Ricci curvature $\beta > 0$.*

Remark 1.1. *In case (1), the Ricci curvature of ω_β, which is $r(\beta) = 1 - \lambda(1-\beta)$ by (1), varies from $r(0) = 1 - \lambda < 0$ to $r(1 - \lambda^{-1} + \epsilon) = \lambda\epsilon > 0$. In particular, there is a Ricci flat conical metric when $\beta = 1 - \lambda^{-1}$, which was also proved by Simon Brendle in [13].*

Our first theorem is

Theorem 1.2. *Assume* $\lambda \geq 1$*. If we already know* X *has a Kähler-Einstein metric, then there exists a conical Kähler-Einstein metric* ω_β *for any* $\beta \in (0,1]$*. Moreover,* ω_β *converges to a unique Kähler-Einstein metric* ω^D_{KE} *as* $\beta \to 1$ *in a weak sense.*

Remark 1.2. *When* $Aut(X)$ *is discrete, we prove more precise result in Theorem 7.1 showing the potential of* ω_β *converge to the potential of* ω_{KE} *in* C^0*-norm. When* $Aut(X)$ *is continuous, the convergence in the last statement is proved modulo one technical point. The point is that, since we need to work in different function space corresponding to different cone angles, the application of implicit function theorem is more delicate as shown by Donaldson in* [20]*. However, in this case, we think the interest of this convergence result is to identify the correct limit Kähler-Einstein metric in the moduli space. For more details, see the discussion in section 7. An algebro-geometric counterpart of this theorem was discussed in* [52]*,* [41]*.*

Using the existence part of above theorem, we can construct a lot of smooth Kähler-Einstein metrics on Fano manifolds using branch covers. More precisely, using the notation of branch-covering before

Theorem 1.3. *If there is conical Kähler-Einstein metric on* $(X, (1-1/m)D)$*, then there is a smooth Kähler-Einstein metric on* Y*. In particular, if* X *admits a Kähler-Einstein metric and* $\lambda \geq 1$*, then there exists smooth Kähler-Einstein metric on* Y*.*

For examples of this construction, see section 4.4.

The proof of existence statement in Theorem 1.2 follows from the following observation. One point is that the log-Mabuchi-energy is well defined on the space of *admissible functions* denoted by $\hat{\mathcal{H}}(\omega)$, which includes all the Kähler potentials of conical Kähler metrics.

Proposition 1.1. *As functionals on* $\hat{\mathcal{H}}(\omega)$*, the log-Mabuchi-energy* $\mathcal{M}_{\omega,(1-\beta)D}$ *is linear in* β*. The log-Ding-energy* $F_{\omega,(1-\beta)D}$ *is concave in* β *up to a bounded constant. As a consequence, consider the interval* $[\beta_1, \beta_2] \subset (0,1]$*. Let* $i \in \{1,2\}$*,* $i^c = \{1,2\} \backslash \{i\}$*. If the log-Mabuchi-energy (resp. log-Ding-energy) is proper for* $\beta = \beta_i$ *and bounded from below for* $\beta = \beta_{i^c}$*, then for any* $\beta_1 < \beta < \beta_2$*, the log-Mabuchi-energy (resp. log-Ding-energy) is proper, so there exists a conical Kähler-Einstein metric on* $(X, (1-\beta)D)$*.*

By combining Proposition 1.1 with the openness result of Donaldson [20], and the result of Berman [6](see section 4.3) we easily see that

Corollary 1.1. *If* $\lambda \geq 1$ *and there is a conical Kähler-Einstein metric on* $((X, (1-\beta)D), c_1(X))$ *for* $0 < \beta < 1$*, then the log-Mabuchi energy* $\mathcal{M}_{\omega,(1-\beta)D}$ *is proper.*

In order to apply this method, we often need to get the lower bound for log-Mabuchi-energy. The following Theorem is very useful.

Theorem 1.4. *If there exists a special degeneration* $(\mathcal{X}, (1-\beta)\mathcal{D}, \mathcal{L})$ *of* $(X, (1-\beta)D)$ *to a conical Kähler-Einstein metric on* $(\mathcal{X}_0, (1-\beta)\mathcal{D}_0)$*. Assume* $\mathcal{X}_0$ *has isolated* $\mathbb{Q}$*-Gorenstein singularities. Then the log-Ding-functional and log-Mabuchi-energy of* $(X, (1-\beta)D)$ *is bounded from below.*

Here the assumption that $\mathcal{X}_0$ has isolated singularities is purely technical, but it is satisfied for our main application here(c.f. Theorem 1.7). One would certainly expect a general statement to be true(see Conjecture 5.1 in Section 5).

Corollary 1.2 (Chen's theorem in the Kähler-Einstein case)**.** *If there exists a special degeneration of Fano manifold* (X, J) *to a Kähler-Einstein manifold* (X_0, J_0)*, then the Mabuchi energy on* X *in the class* $c_1(X)$ *is bounded from below.*

The above results deal with existence question, one may ask for the obstruction of conical Kähler-Einstein metrics. In this direction, we have the following

Theorem 1.5. *If the log-Ding-functional* $F_{\omega,(1-\beta)D}$ *or the log-Mabuchi-energy* $\mathcal{M}_{\omega,(1-\beta)D}$ *for the pair is bounded from below(resp. proper), then the polarized pair* $((X, (1-\beta)D), -K_X)$ *is log-K-semistable(resp. log-K-stable).*

Corollary 1.3.
(1) *If there exists a conical Kähler-Einstein metric on* $((X,(1-\beta)D),c_1(X))$*, then* $((X,(1-\beta)D),-K_X)$ *is log-K-semistable. As a consequence, if* $\lambda>1$*, then* $((X,(1-\beta)D),-K_X)$ *is log-K-semistable for* $0\le\beta<1-\lambda^{-1}+\epsilon$ *for* $0<\epsilon\ll 1$*. If* $\lambda=1$*, then* $(X,(1-\beta)D)$ *is log-K-semistable for* $0\le\beta<\epsilon$*.*
(2) *Assume* $\lambda\ge 1$ *and* $0<\beta<1$*. If there exists a conical Kähler-Einstein metric on* $((X,(1-\beta)D),c_1(X))$*, then* $((X,(1-\beta)D),-K_X)$ *is log-K-stable.*

However, on log-Fano manifold, the cone angle can't be very small. We have

Theorem 1.6. *If* $\lambda<1$*, then* $(X,(1-\beta)D)$ *is not log-K-stable if* $\beta<(\lambda^{-1}-1)/n$*.*

As an illustration of the combination of our methods, we prove

Theorem 1.7. *If* D *is a smooth conic on the projective plane. Then there exists a conical Kähler-Einstein metric on* $(\mathbb{P}^2,(1-\beta)D)$ *if and only if* $\frac{1}{4}<\beta\le 1$*.*

The proof is by constructing a special degeneration to the conic Kähler-Einstein pair. Then we apply Theorem 1.4 and the interpolation property to finish the proof. From the proof we may also speculate the limit of the family of conical Kähler-Einstein metrics when β tends to $1/4$, see conjecture 3 in section 6.

As an application, Theorem 1.7 implies

Corollary 1.4. *A three dimensional* A_2 *singularity* $x_1^2+x_2^2+x_3^2+x_4^3=0$ *admits a Calabi-Yau cone metric with the natural Reeb vector field.*

This settles a question in [24]. Note that however, this contradicts the result by Conti in [17]. We will leave this incompatibility for the study in future.

The organization of the paper is as follows. In section 2, we review some preliminary materials, including Donaldson's Schauder estimate for conical Kähler metrics, various functionals appearing in the theory of Kähler-Einstein metric problem and some existence results for geodesic equation in the space of Kähler metrics. In Section 3, we explain the obstructions to the existence of conical Kähler-Einstein metrics. In particular, we prove Theorem 1.5 and its corollary 1.3. As a particular case, we prove Theorem 1.6. In Section 4, we first review the recent existence result of conical Kähler-Einstein metrics based on the work of Donaldson [20], Jeffres-Mazzeo-Rubinstein-Li [31] and Berman [6]. Then we show our main observation Proposition 1.1, and then prove Theorem 1.3 and existence part of Theorem 1.2, which allow us to construct many smooth Kähler-Einstein metrics using branch covers. In section 5, we prove Theorem 1.4. In Section 6, by combining our method, we prove Theorem 1.7 which shows that the conical Kähler-Einstein metric on $\mathbb{P}^2$ along a smooth conical can only have cone angle in the interval $(\pi/2,2\pi]$. In section 7, we finish the proof of Theorem 1.2 by showing that on a Kähler-Einstein Fano manifold, the conical Kähler-Einstein metrics converge to a unique smooth Kähler-Einstein metric. This is done by carrying out bifurcation analysis first used by Bando-Mabuchi.

After finishing the draft of this paper, we received the paper by Jiang Song and Xiaowei Wang [51]. In the last section 8, we discuss the relation of their work to our paper. Their results have some overlaps with ours. But mostly importantly from our point of view, they proved an existence result in the toric case. The conic Kähler-Einstein spaces they obtained can serve as the degeneration limits of toric Fano manifolds with some smooth pluri-anticanonical divisors. So combining their existence result in the toric case with the strategy in this paper, we show, in the toric case, a version of Donaldson's conjecture which relates the maximal cone angle and the greatest lower bound of Ricci curvature. To state this result, first define

$$R(X)=\sup\{t|\exists\ \omega\in c_1(X)\text{ such that }Ric(\omega)\ge t\omega\}. \tag{2}$$

Proposition 1.2. *For each λ sufficiently divisible, there exists a sub-linear system $\mathscr{L}_\lambda$ of $|-\lambda K_X|$ such that for any general member $D \in \mathscr{L}_\lambda$, if D is smooth, then there exists a conical Kähler-Einstein metric on $(X, (1-\gamma)\lambda^{-1}D)$ if and only if $\gamma \in (0, R(X))$.*

Remark 1.3. *The smoothness assumption is easily satisfied when* $\dim(\mathrm{X}) \leq 2$. *It seems to be guaranteed by choosing $\mathscr{L}_\lambda$ more carefully. See the discussion in Remark 8.2. In general, if D is not smooth, then there exists a weak solution (i.e. bounded solution) to the conical Kähler-Einstein metric.*

The idea of the proof is similar to the proof of Theorem 1.7. The sublinear system $\mathscr{L}_\lambda$ we construct has the property that for each general member D in $\mathscr{L}_\lambda$, (X, D) has a degeneration to the toric conic Kähler-Einstein pair (X, D_0) constructed by Song-Wang.

The interpolation properties of energy functionals obtained in this paper seem to be known to some other experts in the field too. In particular, we were informed by Professor Arezzo that he also observed this.

Acknowledgement: We are grateful to Professor Jian Song and Xiaowei Wang for sending their paper to us. Their existence result in the toric case is a main impetus for us to write the last section in this paper. We also thank them for pointing out our incorrect argument showing that generic divisor appearing in the proof of Proposition 1.2 is smooth. We are indebted to Dr. H-J. Hein for pointing out Corollary 1.4, and related interesting discussions. We thank Professor Xiuxiong Chen, Professor S.K. Donaldson, Professor Gang Tian for encouraging discussions. Part of this work was done during Chi Li's visit to Imperial College and University of Cambridge in UK. He would like to thank Professor S.K.Donaldson and Professor Julius Ross for the invitation which made this joint work possible.

2. PRELIMINARIES

2.1. Space of admissible potentials for conical Kähler metrics.

$$PSH_\infty(\omega) := \{\phi \in L^\infty(X); \omega + \sqrt{-1}\partial\bar{\partial}\phi \geq 0 \text{ and } \phi \text{ is u.s.c. }\}.$$

Definition 2.1. (1) *A conical Kähler metric on $((X, (1-\beta)D), c_1(X))$ is a Kähler current ω in the class $c_1(X)$ such that*
  - *locally, we can choose $\phi \in L^\infty_{loc}$ such that $\omega = \sqrt{-1}\partial\bar{\partial}\phi$.*
  - *ω is smooth on $X \backslash D$*
  - *For any point $p \in D$, we choose local coordinate $\{z_i\}$ in a neighborhood of p such that $D = \{z_1 = 0\}$. Then ω is quasi-isometric to following model metric:*

$$\frac{dz_1 \wedge d\bar{z}_1}{|z_1|^{2(1-\beta)}} + \sum_{i=2}^n dz_i \wedge d\bar{z}_i$$

Geometrically, ω represents a Kähler metric with cone singularities along D of cone angle $2\pi\beta$. In this paper, for simplicity, we will call such metric a conical Kähler metric on $(X, (1-\beta)D)$, i.e. omitting writing the class $c_1(X)$.

(2) *A conical Kähler-Einstein metric on $((X, (1-\beta)D), c_1(X))$ is a conical Kähler metric solving the following equation*

$$Ric(\omega) = r(\beta)\omega + (1-\beta)\{D\}. \tag{3}$$

Here $\{D\}$ is the current of integration on D, and $r(\beta) = 1 - (1-\beta)\lambda$.

Again, we will ommit the class $c_1(X)$ in this paper for simplicity.

Example 2.1. *If $\omega_{sm} \in c_1(X)$ is a smooth Kähler form. Let $[D]$ be the line bundle induced by the divisor D. Choose any Hermitian metric h on $[D]$. Assume $s_D \in H^0(X, [D])$ is the defining section D, i.e. $D = \{s_D = 0\}$. We let*

$$\omega = \omega_{sm} + \epsilon\sqrt{-1}\partial\bar{\partial}|s|_h^{2\beta}$$

It's easy to calculate that

$$\omega = (1-\epsilon\lambda|s|^{2\beta})\omega_{sm} + \epsilon\beta^2|s|_h^{2\beta}\frac{(\nabla s,\overline{\nabla s})_h}{|s|_h^2}$$

So if ϵ *is sufficiently small, this is a positive definite form. For any point* $p \in D$*, if we choose local coordinate chart* $\{z_i\}$ *and local trivializtion* e *such that* $s = z_1 e$ *and* $|s|_h^2 = a|z_1|^2$*, then*

$$\omega = (1-\epsilon\lambda|s|^{2\beta})\omega_{sm} + \epsilon\beta^2 a^\beta|z_1|^{2\beta}\frac{(dz_1+z_1a^{-1}\partial a)\wedge(d\bar z_1+\bar z_1 a^{-1}\bar\partial a)}{|z_1|^2}$$

In order to carry out analysis for the conical metric , one needs to set up good function space such that differential operators behaves well in such space. This is carried out by Donaldson in [20]. We first write the metric and differential operator in the conical coordinate. Let $z_1 = re^{i\theta}$ be the ordinary complex coordinate. Define $\rho = r^\beta$. The model case tells us the right thing to do.

$$\begin{aligned}\omega &= \frac{dz_1\wedge d\bar z_1}{|z_1|^{2(1-\beta)}} + \sum_{j>1} dz_j\wedge d\bar z_j = \frac{(dr+ird\theta)\wedge(dr-ird\theta)}{r^{2(1-\beta)}} + \sum_{j>1} dz_j\wedge d\bar z_j \\ &= (d\rho + i\beta\rho d\theta)\wedge(d\rho - i\beta\rho d\theta) + \sum_{j>1} dz_j\wedge d\bar z_j\end{aligned}$$

Let $\epsilon = e^{i\beta\theta}(d\rho + i\beta\rho d\theta) = d\zeta$, where $\zeta = z_1^\beta = \rho e^{i\beta\theta}$ is defined by choosing a branch locally away from 0. The general (1,1) form, in particular the Kähler form for a conical metric, also has cross terms:

$$\omega = \sqrt{-1}\left(f\epsilon\wedge\bar\epsilon + f_{\bar j}\epsilon\wedge d\bar z_j + f_j dz_j\wedge\bar\epsilon + f_{i\bar j}dz_i\wedge d\bar z_j\right) \tag{4}$$

If $\omega = \sqrt{-1}\partial\bar\partial\phi$,

$$f = \frac{\partial^2\phi}{\partial\zeta\partial\bar\zeta} = \frac{1}{4}\Delta_{\mathbb{C}^2,\beta}\phi = \frac{1}{4}\left(\frac{\partial^2}{\partial\rho^2} + \frac{1}{\rho}\frac{\partial}{\partial\rho} + \frac{1}{\beta^2\rho^2}\frac{\partial^2}{\partial\theta^2}\right)\phi$$

where $\Delta_{\mathbb{C}^2,\beta}$ is the Laplacian associated with the standard conical metric on $\mathbb{C}^2$: $g_\beta = d\rho^2 + \beta^2\rho^2 d\theta^2$.

$$f_{\bar j} = \frac{\partial^2\phi}{\partial\zeta\partial\bar z_j} = \frac{1}{2}e^{-i\beta\theta}\left(\frac{\partial}{\partial\rho} - \frac{i}{\beta\rho}\frac{\partial}{\partial\theta}\right)\frac{\partial}{\partial\bar z_j}\phi$$

Note that in example 2.1, one can take $\phi = f + \epsilon a^\beta|z_1|^{2\beta} = f + \epsilon a^\beta|\zeta|^2$, where f and a are smooth functions of $\{z_i\}$. Since locally $dz = d\zeta^{1/\beta} = \beta^{-1}\zeta^{\frac{1}{\beta}-1}d\zeta$, $\partial\phi = \alpha + \epsilon a^\beta\bar\zeta d\zeta$ with $\alpha = O(1)$ as $z_1 \to 0$. So for ω in the representation (4), $f_{\bar j} = \epsilon\beta a^{\beta-1}a_{\bar j}\bar\zeta = O(|\zeta|)$. We certainly want such ϕ to be in our space. Following Donaldson, we define

Definition 2.2. (1) *f is in $C^{,\gamma,\beta}$ if f is γ-Hölder continuous in the coordinate*

$$(\hat\zeta = \rho e^{i\theta} = r^\beta e^{i\theta} = z|z|^{\beta-1}, z_j)$$

(2) *A (1,0)-form $\alpha = f_1\epsilon + \sum_{j>1} f_j dz_j$ is in $C^{,\gamma,\beta}$ if $f_i \in C^{,\gamma,\beta}$ for $1 \le i \le n$, and $f_1 \to 0$ as $z_1 \to 0$.*

(3) *A (1,1)-form ω written as (4) is in $C^{,\gamma,\beta}$ if $f, f_j, f_{\bar j}, f_{i\bar j} \in C^{,\gamma,\beta}$, and $f_j, f_{\bar j} \to 0$ as $z_1 \to 0$.*

(4) *A function f is in $C^{2,\gamma,\beta}$ if $f, \partial f, \partial\bar\partial f$ are all $C^{,\gamma,\beta}$.*

Remark 2.1. *The point is the derivatives in defining $C^{2,\gamma,\beta}$ function involve only the following wedge differentials:*

$$\frac{\partial}{\partial\rho}, \frac{1}{\rho}\frac{\partial}{\partial\theta}, \frac{\partial}{\partial z_j}, \frac{\partial^2}{\partial\zeta\partial\bar\zeta}, \frac{\partial^2}{\partial\rho\partial\bar z_j}, \frac{1}{\rho}\frac{\partial^2}{\partial\theta\partial\bar z_j}, \frac{\partial^2}{\partial z_i\partial\bar z_j}, \; 2 \le i,j \le n.$$

Donaldson set up the linear theory in [20].

Proposition 2.1. *([20]) If $\gamma < \mu = \beta^{-1} - 1$, the inclusion $C^{2,\gamma,\beta} \to C^{,\gamma,\beta}$ is compact. If ω is a $C^{,\gamma,\beta}$ Kähler metric on (X, D) then the Laplacian of ω defines a Fredholm map $\Delta_\omega : C^{2,\gamma,\beta} \to C^{,\gamma,\beta}$.*

In order to consider the conical Kähler-Einstein metrics for different cone angles at the same time, we define:

Definition 2.3. (1) *(space of admissible functions)*

$$\hat{\mathcal{C}}(X, D) = C^{2,\delta}(X) \cup \bigcup_{0<\beta<1} \left(\bigcup_{0<\gamma<\beta^{-1}-1} C^{2,\gamma,\beta}(X, D) \right)$$

For any $C^{,\gamma,\beta}$ conical metric ω, define

$$\hat{\mathcal{H}}(\omega) = \{\phi \in \hat{\mathcal{C}}(X, D); \omega + \sqrt{-1}\partial\bar{\partial}\phi > 0\}.$$

We will need the following basic openness theorem proved by Donaldson.

Theorem 2.1 ([20]). *Let $\beta_0 \in (0, 1)$, $\alpha < \mu_0 = \beta_0^{-1} - 1$ and suppose there is a smooth conical Kähler-Einstein on $(X, (1 - \beta_0)D)$. If there are no nonzero holomorphic vector fields on X which are tangent to D then for β suifficiently close to β_0 there is a conical Kähler-Einstein on $(X, (1 - \beta)D)$.*

2.2. **Functional from smooth Kähler-Einstein.** In the study of smooth Kähler-Einstein metric problem from variational point of view, we have several energy functionals to consider. These functionals are defined on the space of smooth Kähler potentials:

$$\mathcal{H}(\omega) = \{\phi \in C^\infty(X); \omega_\phi := \omega + \sqrt{-1}\partial\bar{\partial}\phi > 0\}.$$

Now we define the basic functionals in the theory

Definition 2.4. *For any $\phi \in \mathcal{H}(\omega)$, we define the functional*

(1)
$$F^0_\omega(\phi) = -\frac{1}{n+1}\sum_{i=0}^{n}\int_X \phi\omega^i_\phi \wedge \omega^{n-i}/n!$$

(2)
$$J_\omega(\phi) = F^0_\omega(\phi) + \int_X \phi\omega^n/n!$$

(3)
$$I_\omega(\omega_\phi) = \int_X \phi(\omega^n - \omega^n_\phi)/n!,$$

Proposition 2.2. *We have the following relations between these functionals:*

(1)
$$\frac{n+1}{n}J_\omega(\phi) \le I_\omega(\phi) \le (n+1)J_\omega(\phi),$$

(2)
$$F^0_\omega(\phi_2) \ge F^0_\omega(\phi_1) - \int_X (\phi_2 - \phi_1)\omega^n_{\phi_1}/n!$$

(3) *If ϕ_t is a smooth path in $\mathcal{H}(\omega)$, then*

$$\frac{d}{dt}F^0_\omega(\phi_t) = -\int_X \dot{\phi}\omega^n_\phi/n!, \tag{5}$$

Remark 2.2. *Item 3 means $F^0_\omega(\phi)$ is the integral of Bott-Chern form. If we let h be the Hermitian metric on K_X^{-1} such that $\omega_h := -\sqrt{-1}\partial\bar{\partial}\log h = \omega$. Denote $h_\phi = he^{-\phi}$. Connect h and h_ϕ by any path $h_t = he^{-\phi_t}$. The corresponding path of curvature forms $\omega_t = \omega_{h_t} = \omega + \sqrt{-1}\partial\bar{\partial}\phi_t$ connects ω and ω_ϕ. The Bott-Chern form is defined by*

$$BC\left(c_1(K_X^{-1})^{n+1}; h, h_\phi\right) = -\int_0^1 dt(n+1)h_t^{-1}\dot{h}_t c_1(K_X^{-1}, h_t)^n = (n+1)\int_0^1 dt\dot{\phi}\omega_t^n$$

So

$$F^0_\omega(\phi) = -\frac{1}{(n+1)!}\int_X BC\left(c_1(K_X^{-1})^{n+1}; h, h_\phi\right)$$

Next we define the energy functional whose Euler-Lagrange equation is the Kähler-Einstein equation. For any smooth Kähler metric $\omega \in c_1(X)$, there exists a smooth function h_ω, which we call Ricci potential of ω, such that

$$Ric(\omega) - \omega = \sqrt{-1}\partial\bar{\partial}h_\omega, \quad \int_X e^{h_\omega}\frac{\omega^n}{n!} = \int_X \frac{\omega^n}{n!}.$$

Equivalently, $\Omega_\omega := e^{h_\omega}\frac{\omega^n}{n!}$ is the unique volume form satisfying

$$Ric(\Omega_\omega) := -\sqrt{-1}\partial\bar{\partial}\log\Omega_\omega = \omega, \quad \int_X \Omega_\omega = \int_X \frac{\omega^n}{n!}.$$

Definition 2.5. (1) *(Ding-Energy)*

$$F_\omega(\phi) = F^0_\omega(\phi) - V\log\left(\frac{1}{V}\int_X e^{h_\omega - \phi}\omega^n/n!\right)$$

(2) *(Mabuchi-Energy)*

$$\mathcal{M}_\omega(\omega_\phi) = -\int_0^1 dt\int_X (S(\omega_{\phi_t}) - n)\dot{\phi}_t\frac{\omega^n_{\phi_t}}{n!} \tag{6}$$

where ϕ_t is any path connecting 0 and ϕ in the space of Kähler potentials $\mathcal{H}(\omega)$.

We collect various relations between these functionals in the following proposition. These are well known. For the proof, see for example [61].

Proposition 2.3. (1) *The Euler-Lagrange equation of $F_\omega(\omega_\phi)$ is the complex Monge-Ampère equation corresponding to Kähler-Einstein metric equation*

$$(\omega + \sqrt{-1}\partial\bar{\partial}\phi)^n = e^{h_\omega - \phi}\omega^n \Longleftrightarrow Ric(\omega_\phi) = \omega_\phi$$

(2) *Mabuchi energy has an explicit formula:*

$$\begin{aligned}\mathcal{M}_\omega(\omega_\phi) &= \int_X \log\frac{\omega^n_\phi}{e^{h_\omega - \phi}\omega^n}\frac{\omega^n_\phi}{n!} + F^0_\omega(\phi) + \int_X h_\omega\omega^n/n! \\ &= \int_X \log\frac{\omega^n_\phi}{e^{h_\omega}\omega^n}\frac{\omega^n_\phi}{n!} - (I_\omega - J_\omega)(\phi) + \int_X h_\omega\omega^n/n! \end{aligned} \tag{7}$$

(3) *Ding-energy and Mabuchi-energy differ by a cycle. Furthermore, Mabuchi-energy is bounded from below by Ding-energy up to a constant.*

$$F_\omega(\omega_\phi) = \mathcal{M}_\omega(\omega_\phi) - \int_X h_\omega\frac{\omega^n}{n!} + \int_X h_{\omega_\phi}\frac{\omega^n_\phi}{n!} \le \mathcal{M}_\omega(\omega_\phi) - \int_X h_\omega\frac{\omega^n}{n!}.$$

These facts can be generalized to the conical setting (See section 4.1).

Definition 2.6. *A functional $F : \mathcal{H}(\omega) \to \mathbb{R}$ is called proper if there is an inequality of the type*

$$F(\omega_\phi) \ge f\left(J_\omega(\omega_\phi)\right), \text{ for any } \omega_\phi \in \mathcal{H}$$

where $f(t) : \mathbb{R}_+ \to \mathbb{R}$ is some monotone increasing function satisfying $\lim_{t\to+\infty} f(t) = +\infty$.

Now we state a fundamental theorem by Tian which gives equivalent criterion for the existence of Kähler-Einstein metric.

Theorem 2.2 ([60])**.** *If $Aut(X,J)$ is discrete. There exists a Kähler-Einstein metric on X if and only if either $F_\omega(\omega_\phi)$ or $\mathcal{M}_\omega(\omega_\phi)$ is proper on $\mathcal{H}(\omega)$.*

The case when $Aut(X,J)$ is not discrete is more subtle. (We thank Professor Gang Tian, Professor Jiang Song and Professor Robert J. Berman for pointing out this point to us). The full general statement is a conjecture by Tian [60]. But for our application, we just need the following result obtained in [47]. It gives a condition for which Tian's argument for proving the properness works through.

Theorem 2.3 ([47])**.** *If $K \subset G$ is a closed subgroup whose centralizer in G denoted by $\mathrm{Centr_K G}$ is finite, then F_ω is linearly proper on K-invariant potentials.*

Given these theorems, the problem is reduced to the test of properness of F_ω or $\mathcal{M}_\omega$. Tian developed a program using finite dimensional approximations. Define

$$\mathcal{B}_k := \left\{ \frac{1}{k} \log \sum_{\alpha=1}^{N_k} |s_\alpha|^2; \{s_\alpha\} \text{ is a basis of } H^0(X, L^k) \right\} \subset [\omega]$$

to be the space of k-th Bergman metrics. The following conjecture of Tian is the analytic version of Yau-Tian-Donaldson conjecture specialized to the Fano case.

Conjecture 2.1 (Tian)**.** *There is a Kähler-Einstein metric on X if and only if for sufficiently large k, ν_ω is proper on $\mathcal{B}_k$.*

The potentials in $\mathcal{B}_k$ has nice properties:

Lemma 2.1 ([45])**.** *For fixed k, there exists a constant $C_k > 0$ such that for any*

$$C_k^{-1} \cdot (I-J)_\omega(\omega_\phi) \le \mathrm{Osc}(\phi) \le C_k \cdot (I-J)_\omega(\omega_\phi)$$

Remark 2.3. *So the properness in the sense of Definition 2.6 is the same as saying that there exists some monotone increasing function f_k as in Definition 2.6 such that:*

$$\nu_\omega(\omega_\phi) \ge f_k(\mathrm{Osc}(\phi))$$

holds for any $\phi \in \mathcal{B}_k$. In particular, $\mathrm{Osc}(\phi) \to +\infty$ would imply $\nu_\omega(\omega_\phi) \to +\infty$.

2.3. Degenerate Complex Monge-Ampère equation on Kähler manifold with boundary.

Let M be a Kähler manifold of dimension $n+1$ with boundary ∂M. We will be interested in solving the Dirichlet problem of complex Monge-Ampère equation on M. Now we have several results for this problem. First we have the following existence of weak solutions

Theorem 2.4. *([10]) Let ω be a nonnegative, smooth (1,1)-form on X. Assume $\phi_i \in \mathcal{PSH}(\omega) \cap C^0(X)$, $i = 0,1$. Then there exists a bounded geodesic Φ_t connecting ϕ_0 and ϕ_1. In other words, there exists a bounded solution of the Dirichlet problem to the following homogeneous complex Monge-Ampère equation on $X \times [0,1] \times S^1$.*

$$\begin{cases} \pi^*\omega + \sqrt{-1}\partial\bar{\partial}\Phi \ge 0, \\ (\pi^*\omega + \sqrt{-1}\partial\bar{\partial}\Phi)^{n+1} = 0, \\ \Phi|_{X\times\{0\}\times S^1} = \phi_0, \Phi|_{X\times\{1\}\times S^1} = \phi_1. \end{cases} \tag{8}$$

For the reader's convenience, we give the proof by Berndtsson.

Proof. Define

$$\mathcal{K} = \{\Psi_t; \pi^*\omega + \sqrt{-1}\partial\bar{\partial}\Psi \ge 0, \lim_{t\to i} \Psi_t \le \phi_i \text{ for } i = 0,1\}$$

Define the Perron-Bremermann envolope as

$$\Phi(x) = \sup\{\Psi(x); \Psi \in \mathcal{K}\}$$

We want to prove Φ is a bounded solution of (8).

We first define $\tilde{\Psi} = max(\phi_0 - ARe(t), \phi_1 + A(Re(t) - 1))$. For A sufficiently big, $\tilde{\Psi} \in \mathcal{K}$. So we can only consider functions in $\mathcal{K}$ which is greater than $\tilde{\Psi}$. Furthermore, since there is an S^1-symmetry, we can only consider S^1 invariant functions in $\mathcal{K}$. For any such function Ψ, it is convex as function of $Re(t)$, and it satisfies

$$\phi_0 - ARe(t) \le \Psi \le \phi_0 + ARe(t), \quad \phi_1 + A(Re(t) - 1) \le \Psi \le \phi_1 - A(Re(t) - 1).$$

So Ψ has the right boundary values uniformly. The upper semicontinuous regularization Φ^* of Φ satisfy the same estimate and is plurisubharmonic. So $\Phi^* \in \mathcal{K}$. So $\Phi = \Phi^*$ is ω-plurisubharmonic. Since Φ^* is ω-maximal, it satisfies the equation (8). □

Remark 2.4. *The existence of $C^{1,1}$-geodesics for smooth Kähler was first proved by Chen in* [15]. *Since we want to deal with mildly singular varieties, we choose to work with just bounded solutions. There are many other important related works to this result. See for example* [5], [28].

The following Theorem is now a well known fact which comes from many people's work. ([14],[27],[16],[12]). We record a version appeared in [12].

Theorem 2.5. *Assume $\Omega_0 > 0$ is a Kähler form on M and $F > 0$ is a smooth, strictly positive function. Consider the Dirichlet problem of complex Monge-Ampère equation*

$$(\Omega_0 + \sqrt{-1}\partial\bar{\partial}\Phi)^{n+1} = F\Omega_0^{n+1}, \quad \Phi|_{\partial M} = \phi \tag{9}$$

If there exists a smooth subsolution, that is, a $\Psi \in C^\infty(M)$ such that $\Omega_0 + \sqrt{-1}\partial\bar{\partial}\Psi > 0$ and $(\Omega_0 + \sqrt{-1}\partial\bar{\partial}\Psi)^{n+1} \ge F\Omega_0^{n+1}$, then (9) *has a unique solution $\Phi \in C^\infty(M)$.*

We also record a result by Phong-Sturm.

Theorem 2.6. *(*[46]*) Assume $\Omega \ge 0$ and there exists a smooth divisor E in the interior of M such that $\Omega > 0$ on $M \backslash E$. Also assume the line bundle $\mathcal{O}(E)$ has a Hermitian metric H, such that $\Omega_\epsilon = \Omega + \epsilon\sqrt{-1}\partial\bar{\partial}\log H > 0$ for ϵ sufficiently small. If there exists a subsolution $\psi \in C^\infty(M)$. Then the following homogeneous complex Monge-Ampère equation has a bounded solution $\Phi \in L^\infty(M)$*

$$(\Omega + \sqrt{-1}\partial\bar{\partial}\Phi)^{n+1} = 0, \quad \Phi|_{\partial M} = \phi. \tag{10}$$

Moreover, $\Phi \in C^{1,\alpha}(M \backslash E)$ for any $0 < \alpha < 1$.

For the reader's convenience, we sketch the proof of Phong-Sturm's theorem here. By subtracting the subsolution ψ, we can assume $\phi = 0$. Then we approximate the degenerate complex Monge-Ampère equation by a family of non-degenerate equations. Let $\Omega_s = \Omega_0 + s\sqrt{-1}\partial\bar{\partial}\log H$. Consider the family of equations:

$$(\Omega_s + \sqrt{-1}\partial\bar{\partial}\Phi_s)^{n+1} = F_s\Omega_s^{n+1}, \quad \Phi_s|_{\partial M} = 0. \tag{11}$$

By the Theorem 2.5, we can solve these equation for $s > 0$. To get the solution of (10), we want to take limit of Φ_s. So we need to establish uniform a priori estimate for Φ_s. For uniform C^0-estimate for Φ_s, first note that

$$\Delta_{\Omega_\epsilon}\Phi_s \ge -tr_{\Omega_\epsilon}\Omega_s = -tr_{\Omega_\epsilon}(\Omega_\epsilon - (\epsilon - s)\sqrt{-1}\partial\bar{\partial}\log H) = -(n+1) - (\epsilon - s)tr_{\Omega^\epsilon}Ric(H) \ge -C_1,$$

with C_1 independent of s. On the other hand we can solve the Dirichlet boundary problem

$$\Delta_{\Omega_\epsilon}\xi = -C_1, \quad \xi|_{\partial M} = 0.$$

Then by maximal principle, we get $\Phi_s \le \xi$. On the other hand, ψ is a subsolution of (11), by comparison principle of complex Monge-Ampère equation, we get that $\Phi_s \ge \psi$. This uniform C^0-estimate actually allows us to construct generalized solution to the homogeneous complex Monge-Ampère equation.

To get more regularity away from the divisor E, let $\Phi_s^\epsilon = \Phi - (\epsilon - s)\log\|\sigma\|^2$, where σ is the defining section of the line bundle $\mathcal{O}(E)$. We can rewrite this as

$$(\Omega_\epsilon + \sqrt{-1}\partial\bar\partial\Phi_s^\epsilon)^{n+1} = \frac{F_s\Omega_s^{n+1}}{\Omega_\epsilon^{n+1}}\Omega_\epsilon^{n+1} = F_s^\epsilon\Omega_\epsilon^{n+1}$$

Now we choose F_s such that $F_s^\epsilon := F_s\Omega_s^{n+1}/\Omega_\epsilon^{n+1}$ is a constant.

Phong-Sturm in [46] proved the following partial-C^1 (adapting Block's estimate), partial C^2-estimate (adapting Yau's estimate):

$$|\nabla\Phi| \le C_1\|\sigma\|^{-\epsilon A_1}, \quad |\Delta\Phi| \le C_2\|\sigma(z)\|^{-\epsilon A_2}$$

with C_i, A_i independent of s.

3. Obstruction to existence: log-K-stability

3.1. **log-Futaki invariant and log-K-(semi)stability.** Fix a smooth Kähler metric $\omega \in c_1(X)$. Assume D is a smooth divisor such that $D \sim_{\mathbb{Q}} -\lambda K_X$ for some $\lambda > 0 \in \mathbb{Q}$. Assume $\mathbb{C}^*$ acts on (X,D) with generating holomorphic vector field v. There exists a potential function $\theta_v \in C^\infty(X)$ satisfying $\sqrt{-1}\bar\partial\theta_v = \iota_v\omega$. The log-Futaki invariant is defined by Donaldson:

Definition 3.1. *([20])*

$$\begin{aligned} F(-K_X,(1-\beta)D;v) &= F(-K_X;v) + (1-\beta)\left(\int_D \theta_v\frac{\omega^{n-1}}{(n-1)!} - \frac{Vol(D)}{Vol(X)}\int_X \theta_v\frac{\omega^n}{n!}\right) \\ &= -\int_X (S(\omega)-n)\theta_v\frac{\omega^n}{n!} + (1-\beta)\left(\int_D \theta_v\frac{\omega^{n-1}}{(n-1)!} - n\lambda\int_X \theta_v\frac{\omega^n}{n!}\right) \end{aligned} \tag{12}$$

Log-Futaki invariant is an obstruction to the existence of conical Kähler-Einstein metrics as explained in [35]. When $\lambda \ge 1$, one can show that there are no nontrivial $\mathbb{C}^*$ action for the pair (X,D). To obtain the obstruction for the existence we define the log-K-stability by generalizing the original definition by Tian and Donaldson.

Definition 3.2. *A test configuration of (X,L), consists of*

1. *a normal scheme $\mathcal{X}$ with a $\mathbb{C}^*$-action;*
2. *a $\mathbb{C}^*$-equivariant line bundle $\mathcal{L} \to \mathcal{X}$*
3. *a flat $\mathbb{C}^*$-equivariant map $\pi : \mathcal{X} \to \mathcal{C}$, where $\mathbb{C}^*$ acts on $\mathbb{C}$ by multiplication in the standard way;*

such that any fibre $X_t = \pi^{-1}(t)$ for $t \neq 0$ is isomorphic to X and (X,L) is isomorphic to $(X_t, \mathcal{L}|_{X_t})$. The test configuration is called normal if the total space $\mathcal{X}$ is normal.

Any test configuration can be equivariantly embedded into $\mathbb{P}^N \times \mathbb{C}^*$ where the $\mathbb{C}^*$ action on $\mathbb{P}^N$ is given by a 1 parameter subgroup of $SL(N+1,\mathbb{C})$. If Y is any subvariety of X, the test configuration of (X,L) also induces a test configuration $(\mathcal{Y}, \mathcal{L}|_{\mathcal{Y}})$ of $(Y, L|_Y)$.

Let d_k, $\tilde d_k$ be the dimensions of $H^0(X,L^k)$, $H^0(Y,L|_Y^k)$, and w_k, $\tilde w_k$ be the weights of $\mathbb{C}^*$ action on $H^0(X_0,\mathcal{L}|_{X_0}^k)$, $H^0(Y_0,\mathcal{L}|_{Y_0}^k)$, respectively. Then we have expansions:

$$d_k = a_0k^n + a_1k^{n-1} + O(k^{n-2}), \quad w_k = b_0k^{n+1} + b_1k^n + O(k^{n-1})$$

$$\tilde d_k = \tilde a_0k^{n-1} + O(k^{n-2}), \quad \tilde w_k = \tilde b_0k^n + O(k^{n-1})$$

If the central fibre X_0 is smooth, we can use equivariant differential forms to calculate the coefficients by [19]. Let ω be a smooth Kähler form in $c_1(L)$, and $\theta_v = \mathcal{L}_v - \nabla_v$, then

$$a_0 = \int_X \frac{\omega^n}{n!} = Vol(X);\ a_1 = \frac{1}{2}\int_X S(\omega)\frac{\omega^n}{n!} \tag{13}$$

$$b_0 = -\int_X \theta_v\frac{\omega^n}{n!};\ b_1 = -\frac{1}{2}\int_X \theta_v S(\omega)\frac{\omega^n}{n!} \tag{14}$$

$$\tilde{a}_0=\int_{Y_0}\frac{\omega^{n-1}}{(n-1)!}=Vol(Y_0);\ \tilde{b}_0=-\int_{Y_0}\theta_v\frac{\omega^{n-1}}{(n-1)!} \tag{15}$$

Comparing (12), (14)-(15), we can define the algebraic log-Futaki invariant of the given test configuration to be

$$\begin{aligned} F(\mathcal{X},\mathcal{Y},\mathcal{L}) &= \frac{2(a_0b_1-a_1b_0)}{a_0}+(-\tilde{b}_0+\frac{\tilde{a}_0}{a_0}b_0) \\ &= \frac{a_0(2b_1-\tilde{b}_0)-b_0(2a_1-\tilde{a}_0)}{a_0} \end{aligned} \tag{16}$$

Definition 3.3. *(X,Y,L) is log-K-semistable along $(\mathcal{X},\mathcal{L})$ if $F(\mathcal{X},\mathcal{Y},\mathcal{L})\le 0$. Otherwise, it's unstable.*

(X,Y,L) is log-K-polystable along test configuration $(\mathcal{X},\mathcal{L})$ if $F(\mathcal{X},\mathcal{Y},\mathcal{L})<0$, or $F(\mathcal{X},\mathcal{Y},\mathcal{L})=0$ and the normalization $(\mathcal{X}^\nu,\mathcal{Y}^\nu,\mathcal{L}^\nu)$ is a product configuration.

(X,Y,L) is log-K-semistable (resp. log-K-polystable) if, for any integer $r>0$, (X,Y,L^r) is log-K-semistable (log-K-polystable) along any test configuration of (X,Y,L^r).

Remark 3.1. *When Y is empty, then definition of log-K-stability becomes the definition of K-stability. ([60], [19])*

3.2. log-Mabuchi-energy and log-Futaki-invariant.

3.2.1. *Integrate log-Futaki-invariant.* We now integrate the log-Futaki invariant to get log-Mabuchi-energy associated with the problem. Define the functional:

$$F^0_{\omega,D}(\phi)=-\int_0^1 dt\int_D\dot{\phi}\frac{\omega_\phi^{n-1}}{(n-1)!}$$

Then from (5) and (6), we immediately get

Proposition 3.1. *For any $\phi\in C^\infty(X)\cap\mathcal{H}(\omega)$, we can define the log-Mabuchi-energy as*

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)=\mathcal{M}_\omega(\omega_\phi)+(1-\beta)\left(-F^0_{\omega,D}(\phi)+\frac{Vol(D)}{Vol(X)}F^0_\omega(\phi)\right), \tag{17}$$

so that, if $\omega_t=\omega+\sqrt{-1}\partial\bar{\partial}\phi_t$ is a sequence of smooth Kähler metrics in $c_1(X)$, then

$$\frac{d}{dt}\mathcal{M}_{\omega,(1-\beta)D}=-\int_X(S(\omega_t)-n)\dot{\phi}_t\frac{\omega_\phi^n}{n!}+(1-\beta)\left(\int_D\dot{\phi}\frac{\omega_\phi^{n-1}}{(n-1)!}-n\lambda\int_X\dot{\phi}\frac{\omega_\phi^n}{n!}\right)$$

Proposition 3.2. *The log-Mabuchi-energy can be written in the following different forms*

$$\begin{aligned} \mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi) &= \mathcal{M}_\omega(\omega_\phi)+\lambda(1-\beta)(I-J)_\omega(\omega_\phi)+(1-\beta)\int_X\log|s_D|^2(\omega_\phi^n-\omega^n)/n! \\ &= \int_X\log\frac{\omega_\phi^n}{\omega^n}\frac{\omega_\phi^n}{n!}-r(\beta)(I_\omega-J_\omega)(\omega_\phi)+\int_X\left(h_\omega-(1-\beta)\log|s|_h^2\right)\frac{\omega^n-\omega_\phi^n}{n!} \end{aligned}$$

Proof. By the Poincaré-Lelong equation $\frac{\sqrt{-1}}{2\pi}\partial\bar{\partial}\log|s|_h^2=-\lambda\omega+\{D\}$. So

$$\begin{aligned} F^0_{\omega,D}(\phi) &= -\int_0^1 dt\int_X\dot{\phi}\left(\frac{\sqrt{-1}}{2\pi}\partial\bar{\partial}\log|s|_h^2+\lambda\omega\right)\frac{\omega_\phi^n}{n!}=-\int_0^1 dt\int_X\frac{1}{2\pi}\log|s|_h^2\frac{d}{dt}\frac{\omega_t^n}{n!}+\lambda\mathcal{J}^\omega_\omega(\omega_\phi) \\ &= -\int_X\frac{1}{2\pi}\log|s|_h^2\frac{\omega_\phi^n-\omega^n}{n!}+\lambda\mathcal{J}^\omega_\omega(\omega_\phi) \end{aligned}$$

Here, for any smooth closed (1,1)-form χ, we define

$$\mathcal{J}^\chi_\omega(\phi)=-\int_0^1 dt\int_X\dot{\phi}\chi\wedge\frac{\omega_\phi^{n-1}}{(n-1)!}$$

By taking derivatives, it's easy to verify that $nF^0_\omega(\phi)+\mathcal{J}^\omega_\omega=(I-J)_\omega(\omega_\phi)$. So

$$\begin{aligned}&\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)=\mathcal{M}_\omega(\omega_\phi)+\frac{(1-\beta)Vol(D)}{Vol(X)}F^0_\omega(\phi)-F^0_{\omega,(1-\beta)D}(\phi)\\ =\;&\mathcal{M}_\omega(\omega_\phi)+(n\lambda)(1-\beta)F^0_\omega(\phi)+(1-\beta)\lambda\mathcal{J}^\omega_\omega(\phi)+(1-\beta)\int_X\frac{1}{2\pi}\log|s_D|^2(\omega^n_\phi-\omega^n)/n!\\ =\;&\mathcal{M}_\omega(\omega_\phi)+\lambda(1-\beta)(I-J)_\omega(\omega_\phi)+(1-\beta)\int_X\frac{1}{2\pi}\log|s_D|^2(\omega^n_\phi-\omega^n)/n!\end{aligned}$$

□

Remark 3.2. *For any smooth Kähler metric $\omega\in c_1(X)$, we define the twisted Ricci potential $H_{\omega,(1-\beta)D}=h_\omega-\frac{1-\beta}{2\pi}\log|s_D|^2$, so that*

$$\sqrt{-1}\partial\bar\partial H_{\omega,(1-\beta)D}=Ric(\omega)-\omega-(1-\beta)(-\lambda\omega+\{D\})=Ric(\omega)-r(\beta)\omega-(1-\beta)\{D\}.$$

Then we can write the log-Mabuchi-energy in another form:

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)=\int_X\log\frac{\omega^n_\phi}{e^{H_{\omega,(1-\beta)D}}\omega^n}\frac{\omega^n_\phi}{n!}+r(\beta)\left(\int_X\phi\frac{\omega^n_\phi}{n!}+F^0_\omega(\phi)\right)+\int_X H_{\omega,(1-\beta)D}\frac{\omega^n}{n!}\tag{18}$$

using the equality $(I_\omega-J_\omega)(\phi)=-\left(\int_X\phi\omega^n_\phi/n!+F^0_\omega(\phi)\right)$.

Remark 3.3. *For simplicity, by abuse of notation, we will often omit the factor $\frac{1}{2\pi}$ in the following calculations.*

3.2.2. *log-Futaki invariant and asymptotic slope of log-Mabuchi-energy.* In this section, we generalize Sean Paul's work in [43] to the conical setting and prove Theorem 1.5 using the argument from [60] and [45].

Assume $X\subset\mathbb{P}^N$ is embedded into the projective space and ω_{FS} is the standard Fubini-Study metric on $\mathbb{P}^N$. For any $\sigma\in SL(N+1,\mathbb{C})$, denote $\omega_\sigma=\sigma^*\omega_{FS}|_X$. Sean Paul proved the following representation of Mabuchi-energy when it is restricted to the space of Bergman metrics.

Theorem 3.1 ([43])**.** *Let $X^n\hookrightarrow\mathbb{P}^N$ be a smooth, linearly normal, complex algebraic variety of degree ≥ 2. Let R_X denote the* ***X-resultant*** *(the Cayley-Chow form of X). Let $\triangle_{X\times\mathbb{P}^{n-1}}$ denote the* ***X-hyperdiscriminant*** *of format (n-1) (the defining polynomial for the dual of $X\times\mathbb{P}^{n-1}$ in the Segre embedding). Then there are continuous norms such that the Mabuchi-energy restricted to the Bergman metrics is given as follows:*

$$(n!)\mathcal{M}_\omega(\omega_\sigma)=\log\frac{\|\sigma\cdot\triangle_{X\times\mathbb{P}^{n-1}}\|^2}{\|\triangle_{X\times\mathbb{P}^{n-1}}\|^2}-\frac{\deg(\triangle_{X\times\mathbb{P}^{n-1}})}{\deg(R_X)}\log\frac{\|\sigma\cdot R_X\|^2}{\|R_X\|^2}\tag{19}$$

The most important ingredient is the following identity proved by Sean Paul:

Theorem 3.2 (Hyper-discriminant part in the K-energy)**.** *([43])*

$$\begin{aligned}\log\frac{\|\sigma\cdot\triangle_{X\times\mathbb{P}^{n-1}}\|^2}{\|\triangle_{X\times\mathbb{P}^{n-1}}\|^2}&=(N+n-1)\int_0^1 dt\int_{(X\times\mathbb{P}^{N+n-1})^\vee}\dot\Phi_\sigma\omega^{N+n-1}_{FS(\mathbb{P}^\vee)}\\ &=\int_0^1 dt\int_X\dot\phi_\sigma(n(n+1)\omega^n_\sigma-nRic(\omega_\sigma)\wedge\omega^{n-1}_\sigma)\end{aligned}$$

The other ingredients are

Lemma 3.1 ([66],[42])**.** *There is a continuous norm on the Chow forms, which satisfies, for any projective variety $X\subset\mathbb{P}^N$,*

$$\log\frac{\|\sigma\cdot R_X\|^2}{\|R_X\|^2}=(n+1)\int_0^1 dt\int_X\dot\phi_\sigma\omega^n_\sigma=-(n+1)!F^0_\omega(\phi_\sigma).\tag{20}$$

Lemma 3.2. *The degree of hyper-discriminant* $\triangle_{X\times\mathbb{P}^{n-1}}$ *and degree of Chow form* R_X *and* R_D *are given by*

- *([43])*

$$\deg(\triangle_{X\times\mathbb{P}^{n-1}})=\deg((X\times\mathbb{P}^{n-1})^\vee)=\int_{X\times\mathbb{P}^{n-1}}c_{2n-1}(J(\mathcal{O}_{X\times\mathbb{P}^{n-1}}(1,1)))=n^2n!Vol(X)$$

  where $J(\mathcal{O}_{X\times\mathbb{P}^{n-1}}(1,1))$ *is the jet bundle of* $\mathcal{O}_{X\times\mathbb{P}^{n-1}}(1,1)=\pi_1^*\mathcal{O}_{\mathbb{P}^N}(1)\otimes\pi_2^*\mathcal{O}_{\mathbb{P}^{n-1}}(1)$

-

$$\deg(R_X)=(n+1)\deg(X,K_X^{-1})=(n+1)!Vol(X),\quad \deg(R_D)=n\deg(D,K_X^{-1})=n!Vol(D)$$

Combining the formulas (17), (19), and (20), we get

Corollary 3.1.

$$\begin{aligned}n!\mathcal{M}_{\omega,(1-\beta)D}(\omega_\sigma) &= \log\frac{\|\sigma\cdot\triangle_{X\times\mathbb{P}^{n-1}}\|^2}{\|\triangle_{X\times\mathbb{P}^{n-1}}\|^2}-\frac{\deg(\triangle_{X\times\mathbb{P}^{n-1}})}{\deg(R_X)}\log\frac{\|\sigma\cdot R_X\|^2}{\|R_X\|^2}\\ &\quad+(1-\beta)\left(\log\frac{\|\sigma\cdot R_D\|^2}{\|R_D\|^2}-\frac{\deg(R_D)}{\deg(R_X)}\log\frac{\|\sigma\cdot R_X\|^2}{\|R_X\|^2}\right)\end{aligned}$$

For any one parameter subgroup $\lambda(t)=t^A\in SL(N+1,\mathbb{C})$. Although the log-Mabuchi-energy is not convex along $\lambda(t)$, the above Corollary says that it is the linear combination of convex functionals. As a consequence, we have the existence of asymptotic slope. Define $\omega_{\lambda(t)}=\lambda(t)^*\omega_{FS}|_X$, and $(\mathcal{X}_0,\mathcal{D}_0)=\lim_{t\to 0}\lambda(t)\cdot(X,D)$ in the Hilbert scheme (which is the central fibre of the induced test configuration introduced in section 3.1). Then combined with [45], we have the following expansion

Proposition 3.3.

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_{\lambda(t)})=(F+a)\log t+O(1)\tag{21}$$

where $F=F(-K_X,(1-\beta)D;\lambda)$ *is the log-Futaki invariant.* $a\geq 0\in\mathbb{Q}$ *is nonnegative and is positive if and only if the central fibre* $\mathcal{X}_0$ *has generically non-reduced fibre.*

Remark 3.4. *In fact, if* $\mathcal{X}_0$ *is irreducible, then by ([60], [45]) one can calculate that* $a=c\cdot(\mathrm{mult}(\mathcal{X}_0)-1)$ *for* $c>0\in\mathbb{Q}$.

Without loss of generality, we assume each homogeneous coordinate Z_i are the eigenvector of $\lambda(t)$ on $H^0(X,\mathcal{O}(1))$ with eigenvalues $\lambda_0=\cdots=\lambda_K<\lambda_{K+1}\leq\cdots\leq\lambda_N$. Let $\omega_{\lambda(t)}=\omega_{FS}+\sqrt{-1}\partial\bar{\partial}\phi_t$. Then

$$\phi_t=\log\frac{\sum_i t^{\lambda_i}|Z_i|^2}{\sum_i|Z_i|^2}\tag{22}$$

There are three possibilities for $\mathcal{X}_0$.

1. (non-degenerate case) $\lim_{t\to 0}\mathrm{Osc}(\phi_t)\to+\infty$. By (22), this is equivalent to $\cap_{i=0}^K\{Z_i=0\}\bigcap X\neq\emptyset$.
2. (degenerate case) $\mathrm{Osc}(\phi_t)\leq C$ for C independent of t. This is equivalent to $\cap_{i=0}^K\{Z_i=0\}\bigcap X=\emptyset$. In this case, $\mathcal{X}_0$ is the image of X under the projection $\mathbb{P}^N\to\mathbb{P}^K$ given by $[Z_0,\dots,Z_N]\mapsto[Z_0,\dots,Z_K,0,\dots,0]$ and there is a morphism from $\Phi:X=\mathcal{X}_{t\neq 0}\to\mathcal{X}_0$ which is the restriction of the projection. There are two possibilities.
    - (a) $\deg(\Phi)>1$. In this case, $\mathcal{X}_0$ is generically non-reduced. So $a>0$ in (21).
      Example: Assume $X^n\subset\mathbb{P}^N$ is in general position. Then the generical linear subspace $\mathbb{L}\cong\mathbb{P}^{N-n-1}$ satisfies $\mathbb{L}\cap X=\emptyset$. Let $\mathbb{M}\cong\mathbb{P}^n$ be a complement of $\mathbb{L}\subset\mathbb{P}^N$. Then the projection of $\Phi:\mathbb{P}^N\backslash\mathbb{L}\to\mathbb{M}$ gives a projection $\Phi:X\to\Phi(X)$ with degree equal to the degree of X.

(b) $\deg(\Phi) = 1$. In this case, $\mathcal{X}_0$ is generically reduced. We have the following fact Example: Assume $X^n \subset \mathbb{P}^N$ is in general position. Assume $K \geq n+1$, then $N - K - 1 \leq N - n - 2$. So the generical linear subspace $\mathbb{L} \cong \mathbb{P}^{N-K-1}$ satisfies $\mathbb{L} \cap X = \emptyset$. Let $\mathbb{M} \cong \mathbb{P}^K$ be a complement of $\mathbb{L} \subset \mathbb{P}^N$. Then the projection of $\Phi : \mathbb{P}^N \backslash \mathbb{L} \to \mathbb{M}$ gives a projection $\Phi : X \to \Phi(X)$ with degree 1.

Proposition 3.4. *As a functional on the space $\mathcal{H}(\omega)$ of smooth Kähler potentials, if $\mathcal{M}_\omega(\omega_\phi)$ is bounded from below (resp. proper), then $(X, -K_X, (1-\beta)D)$ is log-K-semistable (resp. log-K-stable).*

Proof. If log-Mabuchi-energy is bounded from below, then $F \leq 0$ by the expansion (21).

Assume $\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)$ is proper on $\mathcal{H}(\omega)$ in the sense of Definition 2.6, then in particular it's proper on $\mathcal{B}_k$. Then by Lemma 2.1 and Remark 2.3 (See [45]), in case 1 or 2(a), $F < 0$.

By [37], we only need to test stability for normal test configuration with central fibre being a klt pair $(\mathcal{X}_0, (1-\beta)\mathcal{D}_0)$. In case 2(b), we have the product test configuration with vanishing Futaki invariant. ☐

Remark 3.5. *Recently, Berman [7] proved general result that a Kähler-Einstein $\mathbb{Q}$-Fano variety is K-polystable. His approach is based on the expansion of Ding-functional along special test configurations which are sufficient to test K-polystability by [37].*

3.3. **log-slope stability and log-Fano manifold.** Recall that when $\lambda < 1$, $r(\beta) = 1 - \lambda(1-\beta)$. So when $\beta = 0$, $r(\beta) = 1 - \lambda > 0$. The metric in this case should correspond to complete metric with infinite diameter and with $Ric = 1 - \lambda > 0$. This contradicts Myers theorem. So we expect when β is very small, there does not exist such conical Kähler-Einstein metrics.

This is indeed the case. To see this, we first generalize Ross-Thomas' slope stability to the log setting. (See [48] [52]). For any subscheme $Z \subset X$, we blow up the ideal sheaf $\mathcal{I}_Z + (t)$ on $X \times \mathbb{C}$ to get the degeneration of X to the deformation to the normal cone $T_Z X$. For the polarization, we denote $\mathcal{L}_c = \pi^* L - cE$, where $E = P_Z(\mathbb{C} \oplus N_Z X)$ is the exceptional divisor, $0 < c <$ Seshadri constant of Z with respect to X.

$$H^0(\mathcal{X}, \mathcal{L}_c^k) = H^0(X \times \mathbb{C}, L^k \otimes ((t) + \mathcal{I}_Z)^{ck}) = \bigoplus_{j=0}^{ck-1} t^j H^0(X, L^k \otimes \mathcal{I}_Z^{ck-j}) \oplus t^{ck} \mathbb{C}[t] H^0(X, L^k)$$

So for k sufficiently large,

$$\begin{aligned} H^0(\mathcal{X}_0, \mathcal{L}_c^k) &= H^0(X, L^k \otimes \mathcal{I}_Z^{ck}) \oplus \bigoplus_{i=0}^{ck} t^j H^0(X, L^k \otimes \mathcal{I}_Z^{ck-j} / \mathcal{I}_Z^{ck-j+1}) \\ &= H^0(X, L^k \otimes \mathcal{I}_Z^{ck}) \oplus \bigoplus_{i=0}^{ck} t^j \frac{H^0(X, L^k \otimes \mathcal{I}_Z^{ck-j})}{H^0(X, L^k \otimes \mathcal{I}_Z^{ck-j+1})} \end{aligned}$$

If

$$\chi(X, L^k \otimes \mathcal{I}_Z^{xk}) = a_0(x) k^n + a_1(x) k^{n-1} + O(k^{n-1})$$

then by the calculation in by Ross-Thomas in [48], we know that

$$b_0 = \int_0^c a_0(x) dx - c a_0, \quad b_1 = \int_0^c (a_1(x) + \frac{1}{2} a_0'(x)) dx - c a_1$$

Similarly,

$$H^0(\mathcal{Y}, \mathcal{L}_c^k) \quad = \quad H^0(X \times \mathbb{C}, L^k \otimes ((t) + \mathcal{I}_Z)^{ck} \otimes \mathcal{O}_X / \mathcal{I}_Y) = H^0(Y \times \mathbb{C}, L^k \otimes ((t) + \mathcal{I}_Z \cdot \mathcal{O}_Y)^{ck})$$

And

$$H^0(\mathcal{Y}_0, \mathcal{L}_c^k) = H^0(Y, L^k \otimes (\mathcal{I}_Z \cdot \mathcal{O}_Y)^{ck}) \oplus \bigoplus_{i=0}^{ck} t^j \frac{H^0(Y, L^k \otimes (\mathcal{I}_Z \cdot \mathcal{O}_Y)^{ck-j})}{H^0(X, L^k \otimes (\mathcal{I}_Z \cdot \mathcal{O}_Y)^{ck-j+1})}$$

So, by [48] again, if

$$\chi(Y, L^k \otimes (\mathcal{I}_Z \cdot \mathcal{O}_Y)^{xk}) = \tilde{a}_0(x) k^{n-1} + O(k^{n-2}),$$

then

$$\tilde{b}_0 = \int_0^c \tilde{a}_0(x)dx - c\tilde{a}_0.$$

So we can calculate the log-Futaki invariant as

$$\begin{aligned} a_0 F(\mathcal{X}, \mathcal{Y}, \mathcal{L}) &= 2(a_0 b_1 - a_1 b_0) + (\tilde{a}_0 b_0 - a_0 \tilde{b}_0) \\ &= a_0(2b_1 - \tilde{b}_0) - b_0(2a_1 - \tilde{a}_0) \\ &= 2a_0 \left(\int_0^c (a_1(x) - \frac{1}{2}\tilde{a}_0(x) + \frac{1}{2}a_0'(x))dx \right) - 2(a_1 - \tilde{a}_0/2) \int_0^c a_0(x)dx. \end{aligned} \tag{23}$$

In other words, we can define the log-slope invariant:

$$\begin{aligned} \mu_c^{\log}((X,Y); \mathcal{I}_Z) &= \frac{\int_0^c (a_1(x) - \frac{1}{2}\tilde{a}_0(x) + \frac{1}{2}a_0'(x))dx}{\int_0^c a_0(x)dx} = \frac{\int_0^c (a_1(x) - \frac{1}{2}\tilde{a}_0(x))dx + \frac{1}{2}(a_0(c) - a_0)}{\int_0^c a_0(x)dx} \\ &= \mu_c(X; \mathcal{I}_Z) - \frac{\int_0^c \tilde{a}_0(x)dx}{2\int_0^c a_0(x)dx}. \end{aligned}$$

$$\mu^{\log}((X,Y)) = \frac{a_1 - \tilde{a}_0/2}{a_0} = -\frac{n}{2} \cdot \frac{(K_X + Y) \cdot L^{n-1}}{L^n} = \mu_c(X) - \frac{nY \cdot L^{n-1}}{2L^n}.$$

Definition 3.4. *We call (X, Y) is log-slope-stable, if for any subscheme $Z \subset X$, we have*

$$\mu_c^{\log}((X,Y); \mathcal{I}_Z) < \mu^{\log}((X,Y))$$

Proposition 3.5. *Let X be a Fano manifold, and D a Cartier divisor which is numerically equivalent to $-\lambda K_X$. Then if $\lambda < 1$, the pair $(X, (1-\beta)D)$ is not log-slope-stable for $\beta < (\lambda^{-1} - 1)/n$. In the log-Fano case, the log-Mabuchi-energy is not bounded from below for very small angle.*

Proof. The idea is to look at the test configuration $\mathcal{X}$ given by deformation to the normal cone to D, as in [52]. By Lemma 3.3, the Seshadri constant of $(-K_X, D)$ is equal to $c = 1/\lambda$. We will calculate the Futaki invariant for the semi test configuration polarized by $\mathcal{L} = L(-\frac{1}{\lambda}D)$ with $L = -K_X$ and show it is negative for $\beta < (\lambda^{-1} - 1)/n$.

In our case, if we choose $Z = D \sim -\lambda K_X$, then the calculation simplifies to

$$a_0(x) = \frac{(L - xD)^n}{n!} = (1 - x\lambda)^n \frac{(-K_X)^n}{n!} = (1 - x\lambda)^n a_0, \quad a_0'(x) = -n\lambda(1 - x\lambda)^{n-1} a_0$$

$$a_1(x) = \frac{-K_X \cdot (L - xD)^{n-1}}{2(n-1)!} = (1 - x\lambda)^{n-1} \frac{(-K_X)^n}{2(n-1)!} = \frac{n}{2}(1 - x\lambda)^{n-1} a_0.$$

We let $Y = (1-\beta)D$, then

$$\tilde{a}_0(x) = \begin{cases} 0 & , \quad \text{when } x > 0 \\ \frac{L^{n-1} \cdot (1-\beta)D}{(n-1)!} = n\lambda(1-\beta)a_0 & , \quad \text{when } x = 0 \end{cases}$$

So

$$\int_0^c a_0(x)dx = \frac{a_0}{\lambda(n+1)}(1 - (1 - c\lambda)^{n+1})$$

$$\int_0^c a_0'(x)dx = a_0(c) - a_0 = a_0((1 - c\lambda)^n - 1).$$

$$\int_0^c a_1(x)dx = \frac{a_0}{2\lambda}(1 - (1 - c\lambda)^n), \quad \int_0^c \tilde{a}_0(x)dx = 0.$$

So the log-Futaki invariant is equal to

$$a_0 F = a_0^2(1-(1-c\lambda)^{n+1})\frac{n}{n+1}\left[(\lambda^{-1}-1)\left(\frac{n+1}{n}\cdot\frac{1-(1-c\lambda)^n}{1-(1-c\lambda)^{n+1}}-1\right)-\beta\right]$$

So we get $F \le 0 \Longleftrightarrow \beta \ge \beta(\lambda, c)$, where

$$\begin{aligned}\beta(\lambda,c) &= (\lambda^{-1}-1)\left(\frac{n+1}{n}\frac{1-(1-c\lambda)^n}{1-(1-c\lambda)^{n+1}}-1\right)\\ &= \frac{\lambda^{-1}-1}{n}\left(1-\frac{n+1}{\sum_{i=0}^{n}(1-c\lambda)^{-i}}\right)\end{aligned}$$

It's easy to see that

$$\beta_{\min} := \sup_{0<c<\lambda^{-1}} \beta(\lambda,c) = \frac{\lambda^{-1}-1}{n}$$

□

Example 3.1. *On $\mathbb{P}^2$, when D is a line, then $(X,(1-\beta)D)$ is unstable for all $\beta \in [0,1)$; when D is a conic, then $(X,(1-\beta)D)$ is unstable for $\beta \in (0,1/4)$, and it will be proved below that it is semi-stable for $\beta = 1/4$, and hence poly-stable for $\beta \in (1/4,1)$. On $\mathbb{P}^1\times\mathbb{P}^1$, when D is a diagonal line, $(X,(1-\beta)D)$ is unstable for $\beta \in (0,1/2)$. By viewing $\mathbb{P}^1\times\mathbb{P}^1$ as a double cover of $\mathbb{P}^2$ along a conic curve (See Remark 6.1 in Section 6.1 for details) we see these observations match. It is an interesting question whether the bounds of β given by the above proposition is sharp for a smooth hypersurface d in $\mathbb{P}^n$ with $d < n+1$.*

Lemma 3.3. *The Seshadri constant of $(-K_X, D)$ is equal to λ^{-1}.*

Proof. Note that $\mathcal{X}_0 = X\cup_{D_\infty} E$. Here $E \cong P(N_D \oplus \mathbb{C})$ is the exceptional divisor and $D \cong D_\infty \subset P(N_D\oplus\mathbb{C})$ is the divisor at infinity. $\mathcal{L}_c|_X = K_X^{-1} - cD = (1-c\lambda)K_X^{-1}$. This is ample if and only if $c < \lambda^{-1}$.

On the otherhand, $\mathcal{L}_c|_{P(N_D\oplus\mathbb{C})} = \pi^* K_X^{-1} + c\mathcal{O}_E(1) = \pi^* K_X^{-1} + cD_\infty$. Let h be a Hermitian metric on $\mathcal{O}(D)$ such that $\omega_h := -\sqrt{-1}\partial\bar\partial\log h$ is a Kähler form. Then if we define $\Omega = \lambda^{-1}\pi^*\omega_h + c\sqrt{-1}\partial\bar\partial\log(1+h)$, this gives a smooth rotationally symmetric (1,1)-form on E. To write Ω in local coordinate, we choose two kinds of coordinate charts on E which covers the neighborhood of zero section D_0 and infinity section D_∞ of $P(N_D\oplus\mathbb{C})$ respectively. To do this, just choose local trivialization of $N_D|_D$ to get holomorphic coordinate along the fibre, which is denoted by ξ. Then $h = a|\xi|^2$ for some smooth positive definite function a. Note that $\omega_h = -\sqrt{-1}\partial\bar\partial\log a$. In this local coordinate one can easily calculate that

$$\Omega = (\lambda^{-1} - c\frac{a|\xi|^2}{1+a|\xi|^2})\omega_h + c\sqrt{-1}\frac{a}{(1+a|\xi|^2)^2}\nabla\xi\wedge\overline{\nabla\xi}.$$

where for simplicity we denote $\nabla\xi = d\xi + \xi a^{-1}\partial a$. For the coordinate at infinity, we use coordinate change $\eta = \xi^{-1}$, then

$$\Omega = (\lambda^{-1} - c\frac{a}{a+|\eta|^2})\omega_h + c\sqrt{-1}\frac{a}{(|\eta|^2+a)^2}\nabla'\eta\wedge\overline{\nabla'\eta}$$

with $\nabla'\eta = d\eta - \eta a^{-1}\partial a$. So we easily sees that Ω is positive definite if an only if $c < \lambda^{-1}$. The lemma clearly follows from the combination of above discussions. □

The following example is for log-Calabi-Yau case ($\lambda = 1$). We have more to say about this in the last section 8.

Example 3.2. *Let $X = Bl_p\mathbb{P}^2$, $D \in |-K_X|$ be a general smooth divisor. Choose $Z = E$ to be the exceptional divisor. If we perform the operation of deformation to the normal cone, the central fibre is given by $\tilde{\mathcal{X}}_0 = X\cup_{E=D_\infty}\mathbb{P}(\mathbb{C}\oplus\mathcal{O}(-1))$. The Seshadri constant equals 2 and the line bundle $\mathcal{L}_2$ contracts X along its fibration direction and the resulting test configuration has central fibre $\mathcal{X}_0 = \mathbb{P}(\mathbb{C}\oplus\mathcal{O}(-1)) \cong X$. The boundary divisor on $\mathcal{X}_0$ is given by $F + 2D_\infty$ where*

F is the fibre over the intersection point $F\cap E\in E=D_\infty$. Denote $L=K_X^{-1}$ and $Y=(1-\beta)D$. Then the calculation specializes to

$$a_0(x)=\frac{(L-xZ)^2}{2}=4-x-\frac{x^2}{2},\quad a_1(x)=\frac{-K_X\cdot(L-xZ)}{2}=4-\frac{x}{2}.$$

$$\tilde{a}_0(x)=\deg(L-xZ)|_Y=(1-\beta)(8-x).$$

Using formula (23), it's easy to calculate the log-Futaki invariant as

$$F(\mathcal{X},\mathcal{Y},\mathcal{L}_2)=8\left(\frac{7\beta}{3}-2\right).$$

This is negative if and only if $\beta<6/7$. This is compatible with the calculation in [20] (See also [35]), where, instead of taking deformation to normal cone, the same test configuration is generated by one parameter subgroup in the torus action.

4. EXISTENCE OF CONICAL KÄHLER-EINSTEIN METRIC

4.1. log-Mabuchi-energy and log-Ding-energy.

First note that

Proposition 4.1. *The functional $I_\omega(\omega_\phi)$, $J_\omega(\omega_\phi)$, $F^0_\omega(\phi)$ are well defined for $C^{,\gamma,\beta}$ metric ω and $\phi\in\hat{\mathcal{H}}_\omega$. Moreover, the relations in the smooth case also hold in the log case.*

The log-Mabuchi-energy is well defined for Kähler current with mild singularities. In particular, it's well defined for conical Kähler metrics.

Definition 4.1. (1) *For any $C^{,\gamma,\beta}$ metric ω in $[\omega]$, define the twisted Ricci potential by equations:*

$$Ric(\omega)-r(\beta)\omega-(1-\beta)\{D\}=\sqrt{-1}\partial\bar\partial H_{\omega,(1-\beta)D},\quad \int_X e^{H_{\omega,(1-\beta)D}}\frac{\omega^n}{n!}=\int_X\frac{\omega^n}{n!}$$

(2) *(log-Mabuchi-energy) For any $\phi\in\hat{\mathcal{H}}_\omega$*

$$\begin{aligned}\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi) &= \int_X\log\frac{\omega_\phi^n}{e^{H_{\omega,(1-\beta)D}}\omega^n}\frac{\omega_\phi^n}{n!}+r(\beta)\left(\int_X\phi\frac{\omega_\phi^n}{n!}+F^0_\omega(\phi)\right)+\int_X H_{\omega,(1-\beta)D}\frac{\omega^n}{n!}\\ &= \int_X\log\frac{\omega_\phi^n}{e^{H_{\omega,(1-\beta)D}}\omega^n}\frac{\omega_\phi^n}{n!}-r(\beta)(I-J)_\omega(\omega_\phi)+\int_X H_{\omega,(1-\beta)D}\frac{\omega^n}{n!}.\end{aligned}$$

(3) *(log-Ding-energy)*

$$F_{\omega,(1-\beta)D}(\omega_\phi)=F^0_\omega(\phi)-\frac{V}{r(\beta)}\log\left(\frac{1}{V}\int_X e^{H_{\omega,(1-\beta)D}-r(\beta)\phi}\omega^n/n!\right).$$

Proof of Proposition 1.1. Rewrite the log-Mabuchi-energy as:

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)=\int_X\log\frac{\omega_\phi^n}{\omega^n}\frac{\omega_\phi^n}{n!}+r(\beta)\left(\int_X\phi\frac{\omega_\phi^n}{n!}+F^0_\omega(\phi)\right)+\int_X H_{\omega,(1-\beta)D}\frac{\omega^n-\omega_\phi^n}{n!}.$$

We see that the definition of log-Mabuchi-energy does not depend on the normalization of the twisted Ricci potential $H_{\omega,(1-\beta)D}$. So linearity of log-Mabuchi-energy follows from the linearity of $r=r(\beta)=1-\lambda(1-\beta)$ in β and the relation $H_{\omega,(1-\beta_t)D}=(1-t)H_{\omega,(1-\beta_0)D}+tH_{\omega,(1-\beta_1)D}+C_t$.

For the log-Ding-energy case, let $\beta_t=(1-t)\beta_0+t\beta_1$. So by Hölder inequality we get

$$\begin{aligned}\int_X e^{H_{\omega,(1-\beta_t)D}-\beta_t\phi}\frac{\omega^n}{n!} &= \int_X\left(e^{H_{\omega,\alpha_0D}-\beta_0\phi}\right)^{1-t}\left(e^{H_{\omega,\alpha_1D}-\beta_1\phi}\right)^t\frac{\omega^n}{n!}\\ &\le \left(\int_X e^{H_{\omega,(1-\beta_0)D}-\beta_0\phi}\frac{\omega^n}{n!}\right)^{1-t}\left(\int_X e^{H_{\omega,(1-\beta_1)D}-\beta_1\phi}\frac{\omega^n}{n!}\right)^t\end{aligned}$$

So the conclusion easily follows by taking log. □

Proposition 4.2. (1) *The Euler-Lagrange equations of log-Mabuchi-energy and log-Ding-energy are the same:*

$$(\omega+\sqrt{-1}\partial\bar\partial\phi)^n=e^{-r(\beta)\phi}e^{H_{\omega,(1-\beta)D}}\omega^n$$

(2) *The log-Mabuchi-energy and log-Ding-energy differ by a cycle.*

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)=r(\beta)F_{\omega,(1-\beta)D}(\omega_\phi)+\int_X H_{\omega,(1-\beta)D}\omega^n/n!-\int_X H_{\omega_\phi,(1-\beta)D}\omega_\phi^n/n!$$

(3) *log-Mabuchi-energy is bounded from below by log-Ding-energy:*

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)\geq r(\beta)F_{\omega,(1-\beta)D}(\omega_\phi)+\int_X H_{\omega,(1-\beta)D}\omega^n/n! \tag{24}$$

The equality holds if and only if ω_ϕ is a conical Kähler-Einstein metric on $(X,(1-\beta)D)$.

(4) *(co-cycle condition) Assume ω_i are $C^{,\gamma,\beta}$ Kähler metrics in $c_1(X)$, for $i=1,2,3$. Then*

$$\mathcal{M}_{\omega_1,(1-\beta)D}(\omega_2)+\mathcal{M}_{\omega_2,(1-\beta)D}(\omega_3)=\mathcal{M}_{\omega_1,(1-\beta)D}(\omega_3)$$

$$F_{\omega_1,(1-\beta)D}(\omega_2)+F_{\omega_2,(1-\beta)D}(\omega_3)=F_{\omega_1,(1-\beta)D}(\omega_3)$$

Proof. Items 2 and 4 easily follows from the formula relating twisted potentials of two Kähler metrics.

$$\begin{aligned}H_{\omega_\phi,(1-\beta)D} &= H_{\omega,(1-\beta)D}+\log\frac{\omega^n}{\omega_\phi^n}-r(\beta)\phi-\log\left(\frac{1}{V}\int_X e^{H_{\omega,(1-\beta)D}-r(\beta)\phi}\omega^n/n!\right)\\ &= -\left(\log\frac{\omega_\phi^n}{e^{H_{\omega,(1-\beta)D}-r(\beta)\phi}\omega^n}+\log\left(\frac{1}{V}\int_X e^{H_{\omega,(1-\beta)D}-r(\beta)\phi}\omega^n/n!\right)\right)\end{aligned}$$

Item 3 follows from from concavity of log, because we can write

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)=\int_X\log\frac{\omega_\phi^n}{e^{H_{\omega,(1-\beta)D}-r(\beta)\phi}\omega^n}\frac{\omega_\phi^n}{n!}+r(\beta)F_\omega^0(\phi)+\int_X H_{\omega,(1-\beta)D}\frac{\omega^n}{n!}.$$

☐

It's easy to compare energy functional associated with different reference metrics:

Proposition 4.3. *Assume $\omega_2=\omega_1+\sqrt{-1}\partial\bar\partial\phi$ with $\phi\in PSH_\infty(\omega_1)$. Then the log-Mabuchi-energy (or log-Ding-energy) is proper with Kähler metric ω_1 if and only if it's proper with ω_2. In particular, we can choose ω_1 to be smooth and ω_2 to be a $C^{,\gamma,\beta}$ conical metric.*

Proof. By the co-cycle relation for log-Mabuchi-energy (or log-Ding-energy), this follows from the following Lemma. ☐

Lemma 4.1. *If log-Mabuchi-energy or log-Ding-energy is proper (resp. bounded from below) on the space of smooth Kähler potentials, then it's proper (resp. bounded from below) on the space of admissible Kähler potentials.*

Proof. For any admissible function $\phi\in\widehat{\mathcal{H}}(\omega)$, we can find a sequence of smooth Kähler potentials ϕ_i, such that

$$\lim_{i\to+\infty}|\phi_i-\phi|_{C^0}=0,\quad \lim_{i\to+\infty}\omega_{\phi_i}^n=\omega_\phi^n.$$

So $\lim_{i\to+\infty}\mathcal{M}_{\omega,(1-\beta)D}(\omega_{\phi_i})=\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)$. Similarly, $\lim_{i\to+\infty}I_\omega(\omega_{\phi_i})=I_\omega(\omega_\phi)$. So the properness inequality is preserved by approximation and taking the limit. ☐

Lemma 4.2. *Let ω_i, $i=1,2$ be two $C^{,\gamma,\beta}$ metric. Assume $\omega_2=\omega_1+\sqrt{-1}\partial\bar\partial\phi$ for $\phi\in PSH_\infty(\omega_1)$. Then the norm defined by J_{ω_1} and J_{ω_2} are equivalent, that is, for any other metric ω_3,*

$$|J_{\omega_1}(\omega_3)-J_{\omega_2}(\omega_3)|\leq C(\omega_1,\omega_2)$$

Proof. Assume $\omega_3 = \omega_2 + \sqrt{-1}\partial\bar{\partial}\psi = \omega_1 + \sqrt{-1}\partial\bar{\partial}(\phi+\psi)$.

$$\begin{aligned} J_{\omega_1}(\omega_3) - J_{\omega_2}(\omega_3) &= F^0_{\omega_1}(\phi+\psi) - F^0_{\omega_1+\sqrt{-1}\partial\bar{\partial}\phi}(\psi) + \int_X (\phi+\psi)\frac{\omega_1^n}{n!} - \int_X \psi\frac{\omega_2^n}{n!} \\ &= F^0_{\omega_1}(\phi) + \int_X \phi\frac{\omega_1^n}{n!} + \int_X \psi(\omega_1^n - \omega_2^n)/n! \\ &= J_{\omega_1}(\omega_2) + \mathbf{E}. \end{aligned} \tag{25}$$

We can estimate error term $\mathbf{E}$ using integration by part:

$$\mathbf{E} = \int_X \psi(\omega_1 - \omega_2)\wedge\left(\sum_{i=0}^{n-1}\omega_1^{n-1-i}\wedge\omega_2^i\right) = \int_X -\phi(\omega_3-\omega_2)\wedge\left(\sum_{i=0}^{n-1}\omega_1^{n-1-i}\wedge\omega_2^i\right)$$

$$|\mathbf{E}| \le \int_X |\phi|\,(\omega_2+\omega_3)\wedge\left(\sum_{i=0}^{n-1}\omega_1^{n-1-i}\wedge\omega_2^i\right) \le 2n\|\phi\|_{L^\infty} n! Vol(X)$$

□

Theorem 4.1. *([10]) If there exists a conical Kähler-Einstein metric $\omega_{KE,\beta}$ on $(X,(1-\beta)D)$, then $\omega_{KE,\beta}$ obtains the minimum of log-Ding-energy $F_{\omega,(1-\beta)D}(\omega_\phi)$*

For the reader's convenience, we sketch the proof.

Proof. For any admissible Kähler potential ϕ, by Theorem 2.4, there exists a bounded geodesic Φ_t connecting ϕ and ϕ_{KE}. Berndtsson proved in [10] that the log-Ding-energy is convex along the geodesic. Since ϕ_{KE} is the critical point of log-Ding-energy, the result follows from convexity. □

Corollary 4.1. *$\omega_{KE,\beta}$ also obtains the minimum of log-Mabuchi-energy $\mathcal{M}_{\omega,(1-\beta)D}$.*

Remark 4.1. *Although we don't need it, we have the following fact*

Theorem 4.2. *([6]) The log-Mabuchi-energy $\mathcal{M}_{\omega,(1-\beta)D}$ is proper if and only if the log-Ding-energy $F_{\omega,(1-\beta)D}$ is proper.*

4.2. **Existence of conical Kähler-Einstein metric.** Let's review the result in [31] to solve the conical Kähler-Einstein equation.

Theorem 4.3. *([31]) If the log-Mabuchi-energy is proper on $C^{2,\gamma,\beta}$, then there exists a conical Kähler-Einstein metric on $(X,(1-\beta)D)$.*

The idea is to use continuity method as in the proof in the smooth case. So we consider a family of equations.

$$(\omega+\sqrt{-1}\partial\bar{\partial}\psi)^n = e^{H_{\omega,(1-\beta)D}-t\psi}\omega^n \tag{26}$$

This is equivalent to the equation

$$Ric(\omega_\psi) = t\omega_\psi + (r(\beta)-t)\omega + (1-\beta)\{s=0\}. \tag{27}$$

- Step 1: Start the continuity method. Let

  $$\mathcal{S} = \{t\in(-\infty, r(\beta)); (26) \text{ is solvable}\}.$$

  Then $\mathcal{S}$ is non-empty. This is achieved by solving the equation (26) when $t \ll 0$ using Newton-Moser iteration method. See [31] for details.
- Step 2: Openness of solution set. This follows from implicit function theorem thanks to the Fredholm linear theory set up by Donaldson in Theorem 2.1.

- Step 3: C^0-estimate. This is the same as in the smooth case. (cf. [4],[60]). We sketch the proof here. First by taking derivative with respect to t on both sides of (26), we get
$$\Delta_t \dot{\psi}_t = -t\dot{\psi} - \psi$$
where $\Delta_t = \Delta_{\omega_{\psi_t}}$. So
$$\frac{d}{dt}(I-J)_\omega(\omega_\psi) = -\int_X \psi\Delta\dot{\psi}\frac{\omega_\psi^n}{n!} = \int_X (-\Delta_t - t)\dot{\psi}(-\Delta_t)\dot{\psi}\frac{\omega_\psi^n}{n!} \geq 0$$
Now using (27), one can calculate that
$$\begin{aligned}\frac{d}{dt}\mathcal{M}_{\omega,(1-\beta)D}(\psi_t) &= -\int_X H_{\omega,(1-\beta)D}\Delta_{\omega_\psi}\dot{\psi}\frac{\omega_\psi^n}{n!} + \int_X r(\beta)\psi\frac{d}{dt}\frac{\omega_\psi^n}{n!}\\ &= -\int_X n(Ric(\omega) - r(\beta)\omega - (1-\beta)\{D\})\dot{\psi}\frac{\omega_\psi^n}{n!} - r(\beta)\frac{d}{dt}(I-J)_\omega(\omega_\psi)\\ &= -\int_X nt(\omega_\psi - \omega)\dot{\psi}\frac{\omega_\psi^n}{n!} - r(\beta)\frac{d}{dt}(I-J)_\omega(\omega_\psi)\\ &= -(r(\beta)-t)\frac{d}{dt}(I-J)_\omega(\omega_\psi) \leq 0\end{aligned}$$
So along the continuity path (27), the log-Mabuchi-energy is decreasing. By properness of log-K-energy, we see that $(I-J)_\omega(\omega_\psi)$ is bounded from above. Now the C^0-estimate follows from the following Proposition.

Proposition 4.4. *([31],[55])*
(1) $\mathrm{Osc}(\psi_t) \leq I_\omega(\omega_{\psi_t}) + C$ *for some constant C independent of t.*
(2) *(Harnack estimate)* $-\inf_X \psi_t \leq n\sup_X \psi_t$.

- Step 3: C^2-estimate. To get the C^2-estimate, we can use the Chern-Lu's inequality and maximal principle. More precisely, let $\Xi = \log tr_{\omega_\psi}\omega - \lambda\psi$. Then
$$\Delta_{\omega_\psi}\Xi \geq (C_1 - \lambda n) + (\lambda - C_2 tr_{\omega_\psi}\omega). \tag{28}$$
Here
$$C_1 = \inf_{p\in X, v\in T_pX}\frac{Ric(\omega_\psi)(v,\bar{v})}{g(v,\bar{v})}, \quad C_2 = \sup_{p\in X, v,w\in T_pX}\frac{R(\omega)(v,\bar{v},w,\bar{w})}{g(v,\bar{v})g(w,\bar{w})}$$
where $Ric(\omega_\psi)$ (res. $R(\omega)$) is the Ricci curvature (resp. curvature operator) of ω_ψ (resp. ω). By the equation (27), $Ric(\omega_{\psi_t}) \geq t\omega_{\psi_t}$, so $C_1 \geq t$. The other crucial point is the bisectional curvature of ω is bounded from above. (cf. Appendix of [31]).

  To use the maximal principle in the conical setting, one can use Jeffres's trick as in [30]. We add the barrier function $\epsilon\|s\|^{2\gamma'}$ for $0 < \gamma' < \gamma$ so that $\Xi + \epsilon\|s\|^{2\gamma'}$ obtains the maximum at $x_0' \notin D$. We then apply the maximal principle to $\Xi + \epsilon\|s\|^{2\gamma'}$ and let $\epsilon \to 0$.
- Step 4: $C^{,2,\gamma,\beta}$-estimate. There is a Krylov-Evans' estimate in the conical setting as developed in [31]. The proof is similar to the smooth case but adapted to the conical(wedge) Hölder space.
- Use the above a priori estimate, we prove the solution set $\mathcal{S}$ is closed. So $\mathcal{S} = (-\infty, 1]$ and the equation (26) is solvable.

4.3. **Alpha-invariant and small cone angles.** In [6] and [31], Tian's alpha invariant [55] was generalized to the conical setting. We will explain this modification.

Definition 4.2 (log alpha-invariant)**.** *Fix a smooth volume form Ω. Define*
$$\begin{aligned}&\alpha(K_X^{-1}, (1-\beta)D)\\ =\ &\max\left\{\alpha > 0; \exists 0 < C_\alpha < +\infty\ s.t.\ \int_X e^{-\alpha(\phi - \sup\phi)}\frac{\Omega}{|s_D|^{2(1-\beta)}n!} \leq C_\alpha\ for\ any\ \phi \in PSH_\infty(X)\right\}.\end{aligned}$$

Now fix a smooth Kähler metric $\omega \in c_1(X)$. For any $\alpha < \alpha(K_X^{-1}, (1-\beta)D)$, using concavity of log function, we can estimate: for any $\phi \in \hat{\mathcal{H}}(\omega) \subset PSH_\infty(\omega)$,

$$\begin{aligned}
\log C_\alpha &\geq \log\left(\frac{1}{V}\int_X e^{-\alpha(\phi-\sup\phi)}\frac{e^{h_\omega}\omega^n}{n!|s_D|^{2(1-\beta)}}\right) = \log\left(\frac{1}{V}\int_X e^{-\alpha(\phi-\sup\phi)-\log\frac{|s_D|^{2(1-\beta)}\omega_\phi^n}{e^{h_\omega}\omega^n}}\frac{\omega_\phi^n}{n!}\right)\\
&\geq -\frac{1}{V}\int_X \log\left(\frac{|s_D|^{2(1-\beta)}\omega_\phi^n}{e^{h_\omega}\omega^n}\right)\frac{\omega_\phi^n}{n!} + \alpha\left(\sup\phi - \frac{1}{V}\int_X \phi\frac{\omega_\phi^n}{n!}\right)\\
&\geq \frac{1}{V}\left(-\int_X \log\frac{|s_D|^{2(1-\beta)}\omega_\phi^n}{e^{h_\omega}\omega^n}\frac{\omega_\phi^n}{n!} + \alpha I_\omega(\omega_\phi)\right).
\end{aligned}$$

So

$$\begin{aligned}
\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi) &\geq \alpha I_\omega(\omega_\phi) - r(\beta)(I-J)_\omega(\omega_\phi) - C_\alpha\\
&\geq \left(\alpha - r(\beta)\frac{n}{n+1}\right) I_\omega(\omega_\phi) - C_\alpha
\end{aligned}$$

So if

$$\alpha(K_X^{-1}, (1-\beta)D) > \frac{n}{n+1}r(\beta) = \frac{n}{n+1}\left(1-\lambda(1-\beta)\right), \tag{29}$$

then log-Mabuchi-energy is proper for smooth reference metric. To estimate the alpha-invariant, we use Berman's estimate:

Proposition 4.5. *([6])*

$$\alpha(K_X^{-1}, (1-\beta)D) = \lambda\alpha(L_D, (1-\beta)D) \geq \lambda\min\{\beta, \alpha(L_D|_D), \alpha(L_D)\} > 0 \tag{30}$$

Corollary 4.2. *When $\lambda \geq 1$, if*

$$0 < \beta < \min\left(1, (1-1/\lambda) + \frac{n+1}{n}\min\{\alpha(L_D|_D), \alpha(L_D)\}\right), \tag{31}$$

then the log-Mabuchi-energy is proper for smooth reference metric. In particular, when $0 < \beta \ll 1$, the log-Mabuchi-energy is proper. When $\lambda < 1$, we need to assume in addition that $\beta > n(\lambda^{-1}-1)$.

Proof. This follows from (29), (30) and the relation

$$\lambda\beta > \frac{n}{n+1}\left(1-\lambda(1-\beta)\right) \Longleftrightarrow \beta > n(\lambda^{-1}-1)$$

This is automatically true if $\lambda \geq 1$ and $\beta > 0$. □

Corollary 4.3 (Berman,[6])**.** *When $\lambda \geq 1$, there is no holomorphic vector field on X tangent to D.*

Proof. If v is the holomorphic vector field tangent to D, then v generate a one-parameter subgroup $\lambda(t)$. Log-Mabuchi-energy is linear along $\sigma^*\omega$ with the slope given by the log-Futaki-invariant. This is in contradiction to Corollary 4.2. □

Remark 4.2. *This corollary was speculated by Donaldson in [20]. This is also proved using pure algebraic geometry in Song-Wang's recent work [51].*

Proof of Theorem 1.2(Existence part). By the discussion above, when $\lambda \geq 1$, i.e. in the log-canonically-polarized and log-Calabi-Yau case, the log-Mabuchi-energy is proper for $0 < \beta \ll 1$. When there is a smooth Kähler-Einstein metric on X, the Mabuchi energy is bounded from below by [4]. So we can use Proposition 1.1 to get the properness of log-Mabuchi-energy for any $0 < \beta < 1$. Now we use Theorem 4.3 to conclude. □

Proof of Corollary 1.1. Assume there exists a conical Kähler-Einstein metric for $0 < \beta = \beta_0 < 1$. Since we assume $\lambda \geq 1$, there is no holomorphic vector field on X fixing D. By Donaldson's implicit function theorem for conical Kähler-Einstein metrics in [20], there exists a conical Kähler-Einstein metric for $\beta = \beta_0 + \epsilon$ when $\epsilon \ll 1$. So the log-Mabuchi-energy is bounded for $\beta = \beta_0 + \epsilon$. Because log-Mabuchi-energy is proper for $0 < \beta \ll 1$. Then we can use interpolation result Proposition 1.1 to conclude the log-Mabuchi-energy is proper for $0 < \beta < \beta_0 + \epsilon$. ☐

4.4. **Construct smooth Kähler-Einstein metrics on branch covers.** If $D = mD_1$, we can construct branch cover X with branch locus D.

$$\begin{tikzcd}
B \arrow[r, phantom, "\subset"] \arrow[d] & Y \arrow[d, "\pi"] \\
D \arrow[r, phantom, "\subset"] & X
\end{tikzcd}$$

The canonical divisors are related by Hurwitz formula:

$$K_Y = \pi^*(K_X + \frac{m-1}{m}D)$$

In our setting, since $D \sim -\lambda K_X$, we get

$$K_Y^{-1} = \left(1 - \frac{m-1}{m}\lambda\right)\pi^* K_X^{-1} = r(1/m)\pi^* K_X^{-1}$$

1. (Positive Ricci) $-(K_X + (1-1/m)D)$ is ample. This is equivalent to $r(1/m) > 0$.
   Example: $X = \mathbb{P}^2$. Define $\deg Y = (K_Y^{-1})^2$.
   - $\deg D = 2$, $m = 2$, $\lambda = 2/3$. $\deg Y = 4$. $Y = \mathbb{P}^1 \times \mathbb{P}^1$.
   - $\deg D = 3$, $m = 3$, $\lambda = 1$. $\deg Y = 3$. Y is a cubic surface.
   - $\deg D = 4$, $m = 2$, $\lambda = 4/3$. $\deg Y = 2$. Y is $Bl_7\mathbb{P}^2$.
2. (Ricci flat) $K_X + (1-1/m)D \sim 0$. This is equivalent to $r(1/m) = 0$.
   Example: $X = \mathbb{P}^2$.
   - $\deg D = 4$, $m = 4$, $\lambda = 4/3$. Y is a K3 surface in $\mathbb{P}^3$.
   - $\deg D = 6$, $m = 2$, $\lambda = 2$. Y is a K3 surface.
3. (Negative Ricci) $K_X + (1-1/m)D$ is ample.
   Example: X is $\mathbb{P}^2$ and D is a general smooth, degree d curve such that $\lambda = d/3$. Choose $m|d$. Except for the cases already listed above, $K_X + (1-1/m)D$ is ample and there exists an orbifold Kähler-Einstein metric with negative Ricci curvature.

Proposition 4.6. *Fix an orbifold metric ω on $(X, (1-1/m)D)$. The branch cover π induces a map from $\mathcal{PSH}(\omega)$ to $\mathcal{PSH}(\pi^*\omega)$ by pulling back. The energy functionals are compatible with this pull back.*

$$F^Y_{\pi^*\omega}(\pi^*\phi) = mF^X_{\omega,(1-1/m)D}(\phi)$$

$$\mathcal{M}^Y_{\pi^*\omega}(\pi^*\phi) = m\mathcal{M}^X_{\omega,(1-1/m)D}(\phi)$$

Similar relation holds for the functionals $F^0_\omega(\phi)$, I and J.

Proof. For any orbifold metric $\omega \in c_1(X)$, there exists h_ω such that

$$Ric(\omega) - r(1/m)\omega - (1-1/m)\{D\} = \sqrt{-1}\partial\bar\partial H_{\omega,(1-1/m)D}$$

$\tilde\omega = \pi^*\omega$ is a smooth Kähler metric in $r(1/m)^{-1}c_1(Y)$ satisfying

$$Ric(\tilde\omega) - r(1/m)\tilde\omega = \sqrt{-1}\partial\bar\partial\pi^* H_{\omega,(1-1/m)D}$$

So $h_{\tilde\omega} = \pi^* H_{\omega,(1-1/m)D}$ and $e^{h_{\tilde\omega}}\tilde\omega^n = \pi^*(e^{H_{\omega,(1-1/m)D}}\omega^n)$.

$$\int_X e^{H_{\omega,(1-\beta)D} - r(1/m)\phi}\omega^n/n! = \frac{1}{m}\int_Y e^{h_{\tilde\omega} - r(1/m)\pi^*\phi}\tilde\omega^n/n!$$

By the formula for the $F^0_\omega(\phi)$, I, J functional in Definition 2.4, the relation stated in the proposition holds. ☐

Proof of Theorem 1.3. We can choose the reference metric ω on X to be orbifold metric. Then the pull back $\tilde{\omega} = \pi^*\omega$ is a smooth Kähler metric on Y. If $\omega_{KE} = \omega + \sqrt{-1}\partial\bar{\partial}\phi_{KE}$ is the conical Kähler-Einstein metric on $(X, (1-1/m)D, c_1(X))$, then $\tilde{\phi}_{KE} = \pi^*\phi_{KE}$ is the bounded solution of the following Monge-Ampère equation on Y.

$$(\tilde{\omega} + \sqrt{-1}\partial\bar{\partial}\tilde{\phi})^n = e^{h_{\tilde{\omega}} - r(1/m)\tilde{\phi}}\tilde{\omega}^n.$$

Because locally, the covering map is given by $z \to z^m$, by asymptotic expansion obtained in [31, Proposition 4.3], one verifies $\tilde{\phi}_{KE}$ is C^2. So by elliptic theory, $\tilde{\phi}_{KE}$ is indeed a smooth solution of Kähler-Einstein equation on Y. Note that $\pi^*\omega_{KE}$ is in the Kähler class $\pi^*c_1(X) = r(1/m)^{-1}c_1(Y)$. ☐

5. SPECIAL DEGENERATION TO KÄHLER-EINSTEIN VARIETIES

5.1. Kähler current on singular varieties.

We will first establish some standard notations following [22].

Definition 5.1 ($\mathbb{Q}$-Fano variety)**.** *A normal variety X is $\mathbb{Q}$-Fano if X is klt and $-K_X$ is an ample $\mathbb{Q}$-Cartier divisor.*

Assume X is an n-dimensional $\mathbb{Q}$-Fano variety. D is a smooth divisor such that $D \cap X^{sing} = \emptyset$. Define the space of Kähler metrics on $\mathbb{Q}$-Fano varieties following [22].

Definition 5.2. *([22])*

1. *(Plurisubharmonic functions) Let X be a normal analytic space of pure dimension n. A plurisubharmonic (psh) function ϕ on X is an upper semi-continuous function on X with values in $\mathbb{R} \cup \{-\infty\}$, which is not locally $-\infty$, and extends to a psh function in some local embedding $X \to \mathbb{C}^N$. The function ϕ is strongly psh (resp. C^0, resp. C^∞) iff it extends to a strongly psh function (resp. C^0, resp. C^∞) in some local embedding.*

   A pluriharmonic function on X is a real valued continuous function f on X such that the following equivalent condition holds:
   - *f is locally the real part of a holomorphic function*
   - *Given a local embedding $X \to \mathbb{C}^N$, f extends locally to a pluriharmonic function on $\mathbb{C}^N$.*
   - *$f|_{X^{reg}}$ is pluriharmonic.*
2. *(semi-Kähler current) A semi-Kähler, resp. Kähler, resp. smooth Kähler, potential on X is a family $(U_i, \phi_i)_{i\in I}$ where (U_i) is an open covering of X and ϕ_i is a psh function, resp. a strongly psh function, resp. a C^∞-smooth strongly psh function, on U_i such that $\phi_i - \phi_j$ is plurisubharmonic on $U_i \cap U_j$.*
3. *A smooth Kähler metric ω_{sm} on X is a $\sim$-equivalence class of smooth Kähler potentials. A semi-Kähler (resp. Kähler) current on X is a $\sim$-equivalence class of semi-Kähler (resp. Kähler) potentials.*

   A semi-Kähler current ω is said to have L^∞_{loc} (resp. C^0, resp. Hölder continuous) potentials iff, there exists a potential $(U_i, \phi_i)_{i\in I}$ for ω, each ϕ_i is L^∞_{loc} (resp. C^0, resp. Hölder continuous).

In this paper, we will only be interested in the class of bounded plurisubharmonic functions. Note that, by the above definition, we can define $\sqrt{-1}\partial\bar{\partial}\phi$ for any plurisubmonic function ϕ by extending to ambient $\mathbb{C}^N$, taking $\sqrt{-1}\partial\bar{\partial}$ and then restricting to X. So we define

$$PSH_\infty(\omega) := \{\phi \in L^\infty(X); \omega + \sqrt{-1}\partial\bar{\partial}\phi \geq 0 \text{ and } \phi \text{ is u.s.c. }\}.$$

Remark 5.1. *Any function $\phi \in PSH_\infty(X, \omega)$ is of finite self-energy in the sense of Definition 1.1 in [22].*

Remark 5.2 (Orbifold metric induces L^∞ Hermitian metrics)**.** *When the Cartier index of K_X divides r, $K_X^{\otimes r}$ is a line bundle. Any orbifold metric induces a Hermitian metric on $K_X^{\otimes r}$ and hence on $K_X^{-\otimes r}$. In fact, for any point x, we can choose local uniformization chart $\tilde{U} \to U \ni x$*

such that $U=\tilde U/G$ for some finite group G and we choose local coordinates $\{\tilde z_i\}$ on $\tilde U$. Define r=order of G. Then the Cartier index of K_X at x divides r. The r-pluri-anticanonical form $\tilde\tau=(\partial_{\tilde z_1}\wedge\cdots\wedge\partial_{\tilde z_n})^{\otimes r}$ on $\tilde U$ is G-invariant, so it induces a local generator τ of $K_X^{-\otimes r}$ downstairs. If we have an orbifold metric which is locally induced by a smooth G-invariant metric $\tilde g$ on $\tilde U$. We just define the Hermitian metric on $K_X^{-\otimes r}$ by requiring $|\tau|^2=|\partial_{\tilde z}|_{\tilde g}^{2r}=\det(\tilde g)^r$.

Example 5.1. *Let $\mathbb{Z}_4$ acts on $\mathbb{C}^2$ by $\xi:(\tilde z_1,\tilde z_2)\mapsto(\xi\tilde z_1,\xi\tilde z_2)$ where $\xi=\exp(2\pi\sqrt{-1}/4)$. Let $X=\mathbb{C}^2/\mathbb{Z}_4$, then X has an isolated singularity of index 2, which is usually denoted by $\frac{1}{4}(1,1)$. We can embed X into $\mathbb{C}^5$ by defining $u_i=\tilde z_1^{4-i}\tilde z_2^{i}$ for $i=0,\dots,4$.*

We can choose the orbifold metric induced by the following smooth metric on $\tilde U=\mathbb{C}^2$:

$$\tilde\omega=\sqrt{-1}\partial\bar\partial(|z_1|^2+|z_1|^4+|z_2|^2)=(1+4|z_1|^2)dz_1\wedge d\bar z_1+dz_2\wedge d\bar z_2$$

Then $\tilde\tau=(\partial_{\tilde z_1}\wedge\partial_{\tilde z_2})^{\otimes 2}$ induces a generator τ of $K_X^{-\otimes 2}$ with $|\tau|_{\tilde g}^2=(1+4|\tilde z_1|^2)^2=(1+4|u_1|^{1/2})^2$.

By the above discussion, we see that the Hermitian metric determined by an orbifold metric is not a smooth Hermitian metric defined in Definition 5.2. However, by the above discussion, its relative potential to a smooth Hermitian metric is uniformly bounded, so there is no technical difficulty in dealing with them as L^∞-Hermitian metrics.

5.2. **Proof of Theorem 1.4.**

Assume $\pi:(\mathcal{X},-K_{\mathcal{X}/\mathbb{C}})\to\mathbb{C}$ is a special degeneration. Assume for simplicity, $\mathcal{X}$ has only finite many isolated singularities $\{p_i\}$. For any $\phi\in C^\infty(X)$, we are going to solve the homogeneous Monge-Ampère equation

$$(\Omega+\sqrt{-1}\partial\bar\partial\Phi)^{n+1}=0,\Phi|_{S^1\times X}=\phi. \tag{32}$$

Proposition 5.1. *There exists bounded solution Φ for* (32). *$\Phi\in C^{1,\alpha}(\mathcal{X}\backslash\{p_i\})$.*

Proof. If $\mathcal{X}$ is smooth, then this follows from Błocki's Theorem (Theorem 2.5). If $\mathcal{X}$ is singular, we can choose a equivariant resolution $\pi:\tilde{\mathcal{X}}\to\mathcal{X}$. Then we solve the equation on $\tilde{\mathcal{X}}$:

$$(\tilde\Omega+\sqrt{-1}\partial\bar\partial\tilde\Phi)^{n+1}=0,\tilde\Phi|_{S^1\times X}=\phi \tag{33}$$

with $\tilde\Omega=\pi^*\Omega$ being a smooth, closed, non-negative form. By the following Proposition, we have smooth subsolution for (33). So by Phong-Sturm's result (Theorem 2.6), we can get bounded solution $\tilde\Phi$ of (33) and, moreover, $\tilde\Phi$ is $C^{1,\alpha}$ on $\tilde X\backslash E$, where E is exceptional divisor. Since $\tilde\Phi$ is constant on the fibre of the resolution, $\tilde\Phi$ descends to a solution Φ of (32). ☐

As pointed out in the above proof, to apply Theorem 2.6, we need to know the existence of subsolutions. Let $\mathcal{X}^*=\mathcal{X}\backslash\mathcal{X}_0$. To construct such subsolution, we first note that there is an equivariant isomorphism

$$\rho:\mathbb{C}^*\times X\cong\mathcal{X}^*\hookrightarrow\mathcal{X}. \tag{34}$$

Let $\triangle=\{w\in\mathbb{C};|w|\le 1$ be the unit disk and $\mathcal{X}_\triangle=\pi^{-1}(\triangle)$. We embed the special test configuration equivariantly into $\mathbb{P}^N\times\mathbb{C}$:

$$\phi_{\mathcal{X}}:(\mathcal{X},-K_{\mathcal{X}/\mathbb{C}})\hookrightarrow\mathbb{C}\times(\mathbb{P}^N,\mathcal{O}_{\mathbb{P}^N}(1)).$$

We get a smooth S^1-invariant Kähler metric on $\mathcal{X}_\triangle$ by pulling back $\Omega=\phi_{\mathcal{X}}^*(\omega_{FS}+\sqrt{-1}dw\wedge d\bar w)$.

Proposition 5.2. *For any smooth Kähler potential ϕ, there exists a smooth S^1-invariant smooth Kähler metric Ω_Ψ on $\mathcal{X}_\triangle$ such that $\rho^*\Omega_\Psi|_{S^1\times X}=\pi_2^*\omega_\phi$. As a consequence, Ψ is a subsolution of the homogeneous Monge-Ampère equation* (32).

Remark 5.3. *Similar result was proved in* [46]. *For the reader's convenience, we give a proof here.*

Proof. Under the isomorphism (34), we can write

$$\pi_2^*\omega_\phi+\sqrt{-1}dw\wedge d\bar w=\Omega+\sqrt{-1}\partial\bar\partial\Psi_0$$

by taking $\Psi_0 = -\log(h_\phi/\rho^*\phi_{\mathcal{X}}^* h_{FS})$. Let $\eta(w)$ be a smooth cut-off function such that $\eta(w) = 1$ for $|w| \leq 1/3$ and $\eta(w) = 0$ for $|w| \geq 2/3$. Now we define a new metric on $\mathbb{C}^* \times X$:

$$\begin{aligned}\Omega + \sqrt{-1}\partial\bar{\partial}\Psi : &= \pi_2^*\omega_\phi + \sqrt{-1}dw \wedge d\bar{w} - \sqrt{-1}\partial\bar{\partial}(\eta(w)\Psi_0) + a\sqrt{-1}dw \wedge d\bar{w} \\ &= \Omega + \sqrt{-1}\partial\bar{\partial}(\Psi_0 - \eta(w)\Psi_0 + a|w|^2).\end{aligned}$$

In other words we let $\Psi = (1-\eta(w))\Psi_0 + a|w|^2 + c$ for some constant c.

We will show when $\mathbb{R} \ni a \gg 1$ is chosen to be big enough, then we get a smooth Kähler metric on $\mathcal{X}_\triangle$ with the required condition.

For $|w| \geq 2/3$, $\Omega_\Psi = \pi^*\omega_\phi + a\sqrt{-1}dw \wedge d\bar{w}$. So Ω_Ψ is a smooth S^1-invariant Kähler metric for $|w| \leq 1/3$ or $|w| \geq 2/3$. When $|w| \leq 1/3$, $\Omega_\Psi = \Omega + a\sqrt{-1}dw \wedge d\bar{w}$. We can use the glue map ρ to get a smooth S^1-invariant Kähler metric on $\pi^{-1}(\{|w| \leq 1/3\})$. We need to consider the behavior of Ω_Ψ at any point $p \in \mathbb{C}^* \times X$ such that $1/3 < |w(p)| < 2/3$.

$$\begin{aligned}\Omega_\Psi &= \pi^*\omega_\phi - \eta\sqrt{-1}\partial\bar{\partial}\Psi_0 - \Psi_0\sqrt{-1}\partial\bar{\partial}\eta - \sqrt{-1}\left(\partial\eta \wedge \bar{\partial}\Psi_0 + \partial\Psi_0 \wedge \bar{\partial}\eta\right) + (a+1)\sqrt{-1}dw \wedge d\bar{w} \\ &\geq (1-\eta)(\omega_\phi + \sqrt{-1}dw \wedge d\bar{w}) + \eta\Omega - \epsilon\sqrt{-1}\partial\Psi_0 \wedge \bar{\partial}\Psi_0 + (a - \epsilon^{-1}|\eta_w|^2)\sqrt{-1}dw \wedge d\bar{w}\end{aligned}$$

Note that the first two terms together are strictly positive definite. Because on $\mathcal{X}_{|w|\geq 1/3} = \pi^{-1}(\{|w| \geq 1/3\})$, Ψ_0 is well defined smooth, we can choose ϵ sufficiently small and a sufficiently big such that this is a positive form on $\mathcal{X}_{|w|\geq 1/3}$. □

Proof of Theorem 1.4. There exists a metric h_Ω on $K_{\mathcal{X}/\mathbb{C}}^{-1}$ such that $\Omega = -\sqrt{-1}\partial\bar{\partial}\log h_\Omega$. h_Ω defines a volume form on each fibre. If we choose local coordinate $\{z_i\}$ on $\mathcal{X}_t$ and denote $\partial_z = \partial_{z_1} \wedge \cdots \wedge \partial z_n$ and $dz = dz_1 \wedge \cdots \wedge dz_n$. Then the volume form is given by

$$dV(h_\Omega|_{\mathcal{X}_t}) = \left|\partial_z \wedge \overline{\partial_z}\right|^2_{h_\Omega|_{\mathcal{X}_t}} dz \wedge \overline{dz}$$

Let $\mathcal{S}$ be the defining section of the divisor $\mathcal{D}$. Fix the Hermitian metric $\|\cdot\|$ on $\mathcal{O}_{\mathcal{X}}(\mathcal{D})$ such that $-\sqrt{-1}\partial\bar{\partial}\log\|\cdot\|^2 = \lambda\Omega$.

Let $\omega_t = \Omega|_{\mathcal{X}_t}$. To prove the lower boundedness of log-Ding-functional $F_{\omega,(1-\beta)D}$, by Lemma 4.1, we only need to consider smooth Kähler potentials. For any smooth potential $\phi \in C^\infty(X)$, we solve the homogeneous complex Monge-Ampère equation (32) to get the geodesic ray Φ. Then consider the function on the base defined by

$$f(t) = F^0_{\omega_t}(\Phi|_{\mathcal{X}_t}) - \frac{V}{r(\beta)}\log\left(\frac{1}{V}\int_{\mathcal{X}_t} e^{-r(\beta)\Phi}\frac{dV(h_\Omega|_{\mathcal{X}_t})}{\|\mathcal{S}\|^{2(1-\beta)}}\right)$$

Claim 5.1. *$f(t)$ satisfies $\Delta f \geq 0$.*

Assuming the claim, let's finish the proof of Theorem 1.4. By maximal principle of subharmonic function, we have

$$F^X_{\omega_1,(1-\beta)D}(\phi) = f(1) = \max_{t\in\partial\triangle} f(t) \geq f(0) = F^{\mathcal{X}_0}_{\omega_0,(1-\beta)\mathcal{D}_0}(\Phi|_{\mathcal{X}_0})$$

Now since by assumption, there exists a conical Kähler-Einstein metric $\widehat{\omega}_{KE} = \omega_0 + \sqrt{-1}\partial\bar{\partial}\widehat{\phi}_{KE}$ on $(\mathcal{X}_0, (1-\beta)\mathcal{D}_0)$. By Berndtsson's Theorem 4.1, we have

$$F_{\omega_0,(1-\beta)\mathcal{D}_0}(\Phi|_{\mathcal{X}_0}) \geq F_{\omega_0,(1-\beta)\mathcal{D}_0}(\widehat{\phi}_{KE})$$

So combining the above two inequality, we indeed get the lower bound of log-Ding-energy:

$$F^X_{\omega_1,(1-\beta)D}(\phi) \geq F^{\mathcal{X}_0}_{\omega_0,(1-\beta)\mathcal{D}_0}(\widehat{\phi}_{KE})$$

Now, to prove the claim, we write $f(t)$ as parts: $f(t) = \mathrm{I} + \mathrm{II}$:

$$\mathrm{I} = F^0_{\omega_t}(\Phi|_{\mathcal{X}_t}) = -\int_X BC(\Omega^{n+1}, (\Omega + \sqrt{-1}\partial\bar{\partial}\Phi)^{n+1}), \quad \mathrm{II} = -V\log\left(\frac{1}{V}\int_{\mathcal{X}_t} e^{-r(\beta)\Phi}\frac{dV(h_\Omega|_{\mathcal{X}_t})}{\|\mathcal{S}\|^{2(1-\beta)}}\right)$$

For part I, we use the property of Bott-Chern form and the geodesic equation to get that,

$$\begin{aligned}\sqrt{-1}\partial\bar{\partial}\mathrm{I} &= -\int_{\mathcal{X}_t}\sqrt{-1}\partial\bar{\partial}BC(\Omega^{n+1},(\Omega+\sqrt{-1}\partial\bar{\partial}\Phi)^{n+1})\\ &= -\int_{\mathcal{X}_t}(\Omega+\sqrt{-1}\partial\bar{\partial}\Phi)^{n+1}-\Omega^{n+1}\\ &= \int_{\mathcal{X}_t}\Omega^{n+1}\geq 0\end{aligned}$$

For part II, we can write locally $1=\partial_z\otimes dz$ in the decomposition $\mathcal{O}_{\mathcal{X}}=-K_{\mathcal{X}/\mathbb{C}}+K_{\mathcal{X}/\mathbb{C}}$. Then we think $1\in\mathcal{O}_{\mathbb{C}}$ is a holomorphic section in $\pi_*\mathcal{O}_{\mathcal{X}}=\mathcal{O}_{\mathbb{C}}$.

$$\mathrm{II}=-\log\|1\|^2_{L^2}$$

where $\|\cdot\|^2_{L^2}$ is the L^2-metric induced by the singular metric $H=h_\Omega e^{-r(\beta)\Phi}/\|\mathcal{S}\|^{2(1-\beta)}$ on $-K_{\mathcal{X}/\mathbb{C}}$. Then the subharmonicity is given by the next proposition. ☐

Proposition 5.3. II *is a subharmonic function of* t.

Proof. First note that

$$\begin{aligned}-\sqrt{-1}\partial\bar{\partial}\log H &= \Omega+(1-\lambda(1-\beta))\sqrt{-1}\partial\bar{\partial}\Phi+(1-\beta)(-\lambda\Omega+\{\mathcal{D}=0\})\\ &= (1-\lambda(1-\beta))\Omega_\Phi+(1-\beta)\lambda\Omega+(1-\beta)\{\mathcal{D}\}\end{aligned}$$

is a positive current. If $\mathcal{X}$ is smooth, then the subharmonicity follows immediately from Berndtsson's result in [10]. In our case, $\mathcal{X}$ has isolated singularities. We can use divisors to cut out this singularity and reduce the problem to smooth fibrations of Stein manifolds. We apply Berndtsson-Paun's argument in [11]. They construct a sequence of smooth fibrations $\pi_k:\mathcal{X}_k\to\mathbb{C}$, such that

(1) π_k is a smooth fibration. Each fibre is a Stein manifold.
(2) As $k\to+\infty$, $\{\mathcal{X}_k\}$ form an exhaustion of $\mathcal{X}$.

Now, as proved by Berndtsson-Paun, II is the limit of $\mathrm{II}_k=-\log\|1\|^2_k$, where $\|\cdot\|_k$ is the L^2-norm on the direct image $(\pi_k)_*(\mathcal{O}_{\mathcal{X}_k})$. By result in [9], $-\log\|1\|^2_k$ is plurisubharmonic. So we again get that II is plurisubharmonic on the disk $\{|w|\leq 1\}$. Note that, since II is rotationally symmetric, this is same as saying that II is a convex function of $t=|w|$. ☐

Remark 5.4. *When the central fiber is smooth, this is a theorem of Chen* [16], *where a more general statement concerning constant scalar curvature Kähler metrics is proved, using the weak convexity of Mabuchi functional on the space of Kähler metrics. It seems difficult to adapt Chen's argument to the singular setting. The advantage here(in the log setting) is to use Ding's functional, which requires much weaker regularity of the geodesics. A fundamental result of Berndtsson* [10] *says that the Ding functional is genuinely geodesically convex. This technique has been demonstrated in* [10], [8] *and* [7].

Remark 5.5. *During the writing of this paper, the paper by Berman* [7] *appeared in which more general results about subharmonicity of Ding-functional in the singular setting was proved. It seems very likely that his argument implies the following result.*

Conjecture 5.1. *Let* $\pi:(\mathcal{X},\mathcal{D})\to\mathbb{C}$ *be a special degeneration for* (X,D). *Suppose the central fiber* $(\mathcal{X}_0,\mathcal{D}_0)$ *admits a singular Kähler-Einstein metric of cone angle* $2\pi\beta$ *along* $\mathcal{D}_0$. *Then the log-Ding functional* $F_{\omega,(1-\beta)D}$ *is bounded below. As a consequence, the log-Mabuchi functional* $\mathcal{M}_{\omega,(1-\beta)D}$ *is also bounded below.*

6. A CASE STUDY: D=DEGREE 2 CURVE ON $X=\mathbb{P}^2$

6.1. **Proof of Theorem 1.7.**

Proof of Theorem 1.7. We first apply Phong-Song-Sturm-Weinkove's properness result in Theorem 2.3 to show that the Ding-energy $F_{\omega}^{\mathbb{P}^2}$ is proper on the space of $SO(3)$-invariant Kähler metrics. For this, we need to show that the centralizer of $SO(3,\mathbb{R})$ in $SU(3)$ is finite. Indeed, if $\gamma \in \mathrm{Centr}_{\mathrm{SO}(3,\mathbb{R})}\mathrm{SU}(3)$, then $\gamma \cdot C$ is an invariant degree 2 curve under $SO(3,\mathbb{R})$. But there is a unique curve invariant under $SO(3,\mathbb{R})$ which is just C itself. So $\gamma \cdot C = C$ and we conclude $\gamma \in SO(3,\mathbb{R})$. Since the center of $SO(3,\mathbb{R})$ is finite, so the conclusion follows.

By the calculation in Example 3.1, we see that $(\mathbb{P}^2, (1-\beta)D)$ is unstable when $0 < \beta < 1/4$, so there is no conical Kähler-Einstein metric for $\beta \in (0, 1/4)$, by Corollary 4.1 and Proposition 3.4.

To prove the properness for all $\beta \in (1/4, 1]$, by Proposition 1.1, we only need to show the lower boundedness of log-Mabuchi-energy when $\beta = 1/4$. To do this, we construct a special degeneration to conical Kähler-Einstein variety and apply Theorem 1.4. The special degeneration comes from deformation to the normal cone. Let $\tilde{\mathcal{X}}$ be the blow up of $\mathbb{P}^2 \times \mathbb{C}$ along $D \times \{0\}$. Choose the line bundle $\mathcal{L}_{3/2} := \pi^* K_{\mathbb{P}^2}^{-1} - 3/2E$ where E is the exceptional divisor. Then $\mathcal{L}_{3/2}$ is semi-ample and the map given by the complete linear system $|k\mathcal{L}_{3/2}|$ for k sufficiently big contracts the $\mathbb{P}^2$ in the central fibre and we get a special test configuration $\mathcal{X}$ with central fibre being the weighted projective space $\mathbb{P}(1,1,4)$. The weighted projective space inherits an orbifold Kähler-Einstein metric from the standard Fubini-Study metric on $\mathbb{P}^2$ by the quotient map $\mathbb{P}^2 = \mathbb{P}(1,1,1) \to \mathbb{P}(1,1,4)$ given by $(Z_0, Z_1, Z_2) \to (Z_0, Z_1, Z_2^4) =: [W_0, W_1, W_2]$. (See example 8.1 in section 8 for a toric explanation) The induced orbifold Kähler-Einstein metric is the same as the conical Kähler-Einstein metric on $\mathbb{P}(1,1,4)$ which has cone singularity along the divisor $[W_2 = 0]$ with cone angle $2\pi/4$. There is one orbifold singularity on $\mathbb{P}(1,1,4)$ of type $\frac{1}{4}(1,1)$ as explained in example 5.1. But this does not cause any difficulty by the discussion in Section 5.1. So by Theorem 1.4, we get that the log-Ding-energy $F_{\omega,3/4D}^{\mathbb{P}^2}$ is bounded from below. So by Proposition 1.1, $F_{\omega,(1-\beta)D}(\phi)$ is proper for $\beta \in (1/4, 1]$ on the space of $SO(3,\mathbb{R})$ invariant conical metrics. So by the existence theorem explained in section 4.2, we get the existence of conical Kähler-Einstein metric on $(\mathbb{P}^2, (1-\beta)D)$ for any $\beta \in (1/4, 1]$. ☐

Remark 6.1. *When $\beta = 1/2$, there exists an orbifold metric on $(\mathbb{P}^2, 1/2D)$ coming from the branched covering map given by*

$$\begin{aligned} p : \mathbb{P}^1 \times \mathbb{P}^1 &\to \mathbb{P}^2 \\ [U_0, U_1], [V_0, V_1] &\mapsto [U_0V_0 + U_1V_1, i(U_0V_1 + U_1V_0), i(U_0V_0 - U_1V_1)] \end{aligned}$$

This is a degree 2 cover branching along the diagonal $\triangle = \{[U_0, U_1], [U_0, U_1]\} \subset \mathbb{P}^1 \times \mathbb{P}^1$. Note that

$$p(\triangle) = D = \{Z_0^2 + Z_1^2 + Z_2^2 = 0\} \subset \mathbb{P}^2.$$

$Aut^0(\mathbb{P}^1 \times \mathbb{P}^1) = PSL(2,\mathbb{C}) \times PSL(2,\mathbb{C})$. $PSL(2,\mathbb{C})$ acts diagonally on $\mathbb{P}^1 \times \mathbb{P}^1$ and we will denote such action by $PSL(2)^{\mathrm{diag}}$. The map induces a morphism between groups:

$$\begin{aligned} \varphi : PSL(2,\mathbb{C})^{\mathrm{diag}} &\longrightarrow PSO(3,\mathbb{C}) \\ \begin{pmatrix} a & b \\ c & d \end{pmatrix} &\mapsto \frac{1}{ad-bc} \begin{pmatrix} (a^2+b^2+c^2+d^2)/2 & -i(ab+cd) & -i(a^2-b^2+c^2-d^2)/2 \\ i(ac+bd) & ad+bc & ac-bd \\ i(a^2+b^2-c^2-d^2)/2 & ab-cd & (a^2-b^2-c^2+d^2)/2 \end{pmatrix} \end{aligned}$$

Note that $\varphi(SU(2)) = SO(3,\mathbb{R})$. Actually this morphism can be seen as the complexification of the covering map from $SU(2)$ to $SO(3,\mathbb{R})$.

$(\mathbb{Z}_2 \times SU(2)^{\mathrm{diag}})$-invariant metrics on $\mathbb{P}^1 \times \mathbb{P}^1$ are one-to-one correspondent to $SO(3,\mathbb{R})$-invariant orbifold metrics on $(\mathbb{P}^2, 1/2D)$. Now if we take $K = PSU(2,\mathbb{C})^{diag}$, then it's easy to see that the centralizer of K in $G = PSU(2,\mathbb{C}) \times PSU(2,\mathbb{C})$ is finite. By Theorem 2.3, the Ding energy $F_{\omega}^{\mathbb{P}^2}$ is proper on $\mathbb{Z}_2 \times PSU(3)^{diag}$ invariant Kähler metrics. By Propositioin 4.6, we haven shown directly that the log-Ding-energy is proper on the space of $SO(3,\mathbb{R})$ invariant conical Kähler metrics on $(\mathbb{P}^2, 1/2D)$.

From the above proof it is tempting to expect that

Conjecture 6.1. *The conical Kähler-Einstein metric* ω_β *on* $\mathbb{P}^2$ *with cone angle* $2\pi\beta$ *along a smooth degree* 2 *curve converge in the Gromov-Hausdorff sense to the standard orbifold Kähler-Einstein metric on* $\mathbb{P}(1,1,4)$ *as* β *tends to* $1/4$.

Actually, more generally, assume there is a special degeneration $(\mathcal{X},\mathcal{Y})$ of the pair (X,Y), such that $(\mathcal{X}_0,\mathcal{Y}_0)$ is a conical Kähler-Einstein pair. Then we expect (X,Y) converges to $(\mathcal{X}_0,\mathcal{Y}_0)$ in Gromov-Hausdorff sense along certain continuity method (either by the classical continuity method by increasing Ricci curvature (cf. [4],[34]), or by changing cone angles(cf. [20]) , or even by log-Kähler-Ricci flow(cf. in [53]). This philosophy is certainly well known to the expects in the field. In particular, this is related to [62] and [21].

Remark 6.2. *In* [32], *smoothable degenerations of* $\mathbb{P}^2$ *are classified. They are given by smoothings of weighted projective planes* $\mathbb{P}^2(a^2,b^2,c^2)$ *where* (a,b,c) *satisfies the Markov equation:* $a^2+b^2+c^2=3abc$. *Different solutions are related by an operation called mutation:* $(a,b,c)\to(a,b,3ab-c)$. *The first several solutions are* $(1,1,1)$, $(1,1,2)$, $(1,5,2)$, $(1,5,13)$, $(29,5,2)$. *The above construction gives first geometric realization of such degeneration corresponding to the mutation* $(1,1,1)\to(1,1,2)$. *We expect there is similar geometric realization of every mutation.*

6.2. **Calabi-Yau cone metrics on three dimensional** A_2 **singularity.**

Through a stimulating discussion with Dr. Hans-Joachim Hein, we learned that Theorem 1.7 has the following application. Recall it was discovered by Gauntlett-Martelli-Sparks-Yau [24] that there may not exist Calabi-Yau cone metrics on certain isolated quasi-homogeneous hypersurface singularities, with the obvious Reeb vector field. In particular, there are two constraints: Bishop obstruction and Lichnerowicz obstruction. As an example, the case of three dimensional A_{k-1} singularities was studied. Recall a three dimensional A_{k-1} singularity is the hypersurface in $\mathbb{C}^4$ defined by the following equation

$$x_1^2+x_2^2+x_3^2+x_4^k=0.$$

There is a standard Reeb vector field ξ_k which generates the $\mathbb{C}^*$ action with weights $(k,k,k,2)$. Let L_k be the Sasaki link of the A_{k-1} singularity. Then the existence of a Calabi-Yau cone metric with Reeb vector field ξ_k is equivalent to the existence of a Sasaki-Einstein metric on L_k. In [24], using the Bishop obstruction, it was proved that L_k admits no Sasaki-Einstein metric for $k>20$, and using Lichnerowicz obstruction this bound was improved to $k>3$. For $k=2$ this is the well-known conifold singularity and there is a homogeneous Sasaki-Einstein metric on the link L_2. For $k=3$ by Matsushima's theorem the possible Sasaki-Einstein metric on L_3 must be invariant under $SO(3;\mathbb{R})$ action, and is of cohomogeneity one. The ordinary differential equation has been written down explicitly in [24], and it is an open question in [24] whether L_3 admits a Sasaki-Einstein metric.

In the language of Sasaki geometry, the above examples L_k are all quasi-regular, meaning that the Reeb vector field ξ_k generates an S^1 action on L_k, and the quotient M_k is a polarized orbifold M_k(in the sense of [49]). The existence of a Sasaki-Einstein metric on L_k is equivalent to the existence of an orbifold Käler-Einstein metric on M_k. In the above concrete cases, the orbifold M_k is the hypersurface in $\mathbb{P}(k,k,k,2)$ defined by the same equation $x_1^2+x_2^2+x_3^2+x_4^k=0$. Note that $\mathbb{P}(k,k,k,2)$ is not well-formed. When $k=2m+1$ is odd, then

$$\begin{array}{rcl}\mathbb{P}(2m+1,2m+1,2m+1,2) & \xrightarrow{\cong} & \mathbb{P}(2m+1,2m+1,2m+1,2(2m+1))=\mathbb{P}(1,1,1,2)\\ [x_1,x_2,x_3,x_4] & \mapsto & [x_1,x_2,x_3,x_4^{2m+1}].\end{array}$$

Similarly, when $k=2m$ is even, then

$$\begin{array}{rcl}\mathbb{P}(2m,2m,2m,2) & \xrightarrow{\cong} & \mathbb{P}(2m,2m,2m,2m)=\mathbb{P}(1,1,1,1)\\ [x_1,x_2,x_3,x_4] & \mapsto & [x_1,x_2,x_3,x_4^{m}].\end{array}$$

So M_k is isomorphic to $\{z_1^2+z_2^2+z_3^2+z_4=0\}\cong\mathbb{P}^2$ when k is odd, and to $\{z_1^2+z_2^2+z_3^2+z_4^2=0\}\cong\mathbb{P}^1\times\mathbb{P}^1$ when k is even. Regarding the non-well-formed orbifold structure it is not hard to see that when k is odd we get $(\mathbb{P}^2,(1-1/k)D)$ and when k is even we get $(\mathbb{P}^1\times\mathbb{P}^1,(1-2/k)\Delta)$.

Thus we see the close relationship between the existence of Sasaki-Einstein metric on L_k and the existence of conical Kähler-Einstein metric on $(\mathbb{P}^2,(1-1/k)D)$. In particular we know there is no Sasaki-Einstein metric on L_k for $k>3$ by Example 3.1,. This is not surprising at all, since by [49] the Lichnerowicz obstruction could be interpreted as slope stability for orbifolds. The new observation here is the case $k=3$ follows from Theorem 1.7. So we know the three dimensional A_2 singularity admits a Calabi-Yau cone metric with the standard Reeb vector field. This is Corollary 1.4.

The corresponding Sasaki-Einstein metric on L_k is invariant under the $SO(3;\mathbb{R})$ action. It would be interesting to find an explicit solution by solving the ODE written in [24]. In [17] cohomogeneity one Sasaki-Einstein five manifolds were classified, but the above result suggests that the classification is incomplete.

7. Convergence of conical KE to smooth KE

In this section, we prove the convergence statement in Theorem 1.2. So we assume there exists smooth Kähler-Einstein metric on X. When $Aut(X)$ is discrete, then ω_{KE} is invariant under $Aut(X)$. In this case, the Mabuchi energy is proper on $\hat{\mathcal{H}}(\omega)$.

Theorem 7.1. *Assume $\omega_\beta=\omega+\sqrt{-1}\partial\bar\partial\phi_\beta$, and $\omega_{KE}=\omega+\sqrt{-1}\partial\bar\partial\phi_{KE}$ then, ϕ_β converges to ϕ_{KE} in C^0-norm. Moreover, ϕ_β converges smoothly on any compact set away from D.*

Proof. By Theorem 1.1, the log-Mabuchi-energy $\mathcal{M}_{\omega,(1-\beta)D}$ is proper on $\hat{\mathcal{H}}$ for $\beta\in(0,1]$. Furthermore, there exists a constant C independent of β such that

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\phi)\geq C_1 I_\omega(\omega_\phi)-C_2. \tag{35}$$

When $\omega_\phi=\omega_\beta$ is the conical Kähler-Einstein metric on $(X,(1-\beta)D)$ then

$$\mathcal{M}_{\omega,(1-\beta)D}(\omega_\beta)\leq\mathcal{M}_{\omega,(1-\beta)D}(\omega)=0.$$

So from (35) , we see that there exists a constant C independent of β such that

$$I_\omega(\omega_\beta)\leq C.$$

Assume $\omega_\beta=\omega+\sqrt{-1}\partial\bar\partial\phi_\beta$. By Proposition 4.4([31]), there exists a constant C independent of β such that

$$Osc(\phi_\beta)\leq I_\omega(\omega_\beta)+C=\int_X\phi(\omega^n-\omega_\beta^n)/n!+C.$$

So $\|\phi_\beta\|_{C^0}$ is uniformly bounded. Now the theorem follows from standard pluripotential theory. For the last statement, we use the same argument as in the in the proof of existence result in section 4.2. First we use Chern-Lu's inequality:

$$\Delta_\beta(\log tr_{\omega_\beta}\omega-C\phi_\beta)\geq(C_1-\lambda n)+(\lambda-C_2tr_{\omega_\beta}\omega).$$

to get C^2-estimate on ϕ_β. Note that it's easy to verify from the calculation in the Appendix of [31] that we can choose the upper bound on bisectional curvature to be independent of β. Then we can use Krylov-Evan's estimate to get uniform higher order estimate on any compact set away from D. The smooth convergence follows from these uniform estimates. □

When $Aut(X)$ is continuous, then $Aut(X)$ is the complexification of $G:=Isom(X,g_{KE})$. By [4], the moduli space of Kähler-Einstein metrics (denoted by $\mathscr{M}_{KE}$) is isomorphic to the symmetric space $G^{\mathbb{C}}/G$. The important step now is to identify the limit ω_{KE}^D. ω_{KE}^D is the critical point of the following functional, which is part of log-Mabuchi-functional.

Lemma 7.1. *Define the functional*

$$\mathscr{F}_{\omega,D}(\omega_\phi)=\lambda(I-J)_\omega(\omega_\phi)+\int_X\log\|s\|_h^2(\omega_\phi^n-\omega^n)/n!$$

where $\lambda\omega=\sqrt{-1}\partial\bar\partial\log\|s\|_h^2$. Then $\mathscr{F}$ satisfies the following properties:

(1)

$$\mathcal{M}_{\omega,D}(\omega_\phi)=\mathcal{M}_\omega(\omega_\phi)+(1-\beta)\mathscr{F}_{\omega,D}(\omega_\phi). \tag{36}$$

(2) *For $\psi\in\mathcal{PSH}_\infty(\omega)$, we have*

- $$\mathscr{F}_{\omega,D}(\omega_\phi)-\mathscr{F}_{\omega_\psi,D}(\omega_\phi)=\mathscr{F}_{\omega,D}(\omega_\psi).$$
- $$\mathscr{F}_\omega(\omega_\psi)=-\mathscr{F}_{\omega_\psi}(\omega).$$

(3) *$\mathscr{F}$ is convex along geodesics of Kähler metrics. .*

Proof. By calculation in (25),

- $$J_\omega(\omega_\phi)-J_{\omega_\psi}(\omega_\phi)=J_\omega(\omega_\psi)+\int_X(\phi-\psi)(\omega^n-\omega_\psi^n)/n!.$$
- $$\begin{aligned}J_{\omega_\psi}(\omega)+J_\omega(\omega_\psi)&=F^0_\omega(\psi)+\int_X\phi\omega^n/n!+F^0_{\omega_\psi}(-\psi)+\int_X(-\psi)\omega^n/n!\\&=\int_X\psi(\omega^n-\omega_\psi^n)/n!.\end{aligned}$$

We can also calculate

- $$\begin{aligned}I_\omega(\omega_\phi)-I_{\omega_\psi}(\omega_\phi)&=\int_X\phi(\omega^n-\omega_\phi^n)/n!-\int_X(\phi-\psi)(\omega_\psi^n-\omega_\phi^n)/n!\\&=\int_X(\phi-\psi)(\omega^n-\omega_\psi^n)/n!+\int_X\psi(\omega^n-\omega_\phi^n)/n!.\end{aligned}$$
- $$I_\omega(\omega_\psi)+I_{\omega_\psi}(\omega)=2\int_\psi(\omega^n-\omega_\psi^n)/n!.$$

So

- $$(I-J)_\omega(\omega_\phi)-(I-J)_{\omega_\psi}(\omega_\phi)=\int_X\psi(\omega^n-\omega_\phi^n)/n!-J_\omega(\omega_\psi).$$
- $$(I-J)_\omega(\omega_\psi)+(I-J)_{\omega_\psi}(\omega)=\int_X\psi(\omega^n-\omega_\psi^n)/n!.$$

On the other hand,

- $$\log\|s\|_h^2(\omega_\phi^n-\omega^n)-\log\|s\|^2_{he^{-\lambda\psi}}(\omega_\phi^n-\omega_\psi^n)=\lambda\psi(\omega_\phi^n-\omega_\psi^n)+\log\|s\|^2(\omega_\psi^n-\omega^n).$$
- $$\log\|s\|_h^2(\omega_\psi^n-\omega^n)+\log\|s\|^2_{he^{-\lambda\psi}}(\omega^n-\omega_\psi^n)=-\lambda\psi(\omega^n-\omega_\psi^n).$$

So we see that

- $$\mathscr{F}_{\omega,D}(\omega_\phi)-\mathscr{F}_{\omega_\psi,D}(\omega_\phi)=\mathscr{F}_{\omega,D}(\omega_\psi).$$
- $$\mathscr{F}_{\omega,D}(\omega_\psi)+\mathscr{F}_{\omega_\psi,D}(\omega)=0.$$

For the second statement, it is well known that $\mathscr{M}$ is a totally geodesic submanifold of the space of smooth Kähler metrics in $c_1(X)$ and $(I-J)_\omega(\omega_\phi)$ is convex on the space of smooth Kähler metrics. Assume $\phi(t)$ is a geodesic, i.e. $\ddot{\phi}-|\nabla\dot{\phi}|^2_{\omega_\phi}=0$.

$$\begin{aligned}\frac{d}{dt}(I-J)_\omega(\omega_\phi) &= -\int_X \phi\Delta_{\omega_\phi}\dot{\phi}\omega_\phi^n/n! = -\int_X \dot{\phi}(\omega_\phi-\omega)\wedge\omega_\phi^{n-1}/(n-1)!\\ &= n\frac{d}{dt}F^0_\omega(\phi)+\int_X \dot{\phi}\omega\wedge\omega_\phi^{n-1}/(n-1)!.\end{aligned}$$

$$\frac{d}{dt}\int_X \log\|s\|_h^2(\omega_\phi^n-\omega^n)/n! = \int_X \log\|s\|^2\Delta_{\omega_\phi}\dot{\phi}\omega_\phi^n/n! = \int_X(-\lambda\omega+\{D\})\dot{\phi}\omega_\phi^{n-1}/(n-1)!.$$

$$\frac{d}{dt}\mathscr{F}_{\omega,D}(\omega_\phi)=\frac{d}{dt}(n\lambda F^0_\omega(\phi)-F^0_{\omega,D}(\phi))$$

Now since $F^0_\omega(\phi)$ is affine along geodesics of Kähler metrics.

$$\begin{aligned}\frac{d^2}{dt^2}\mathscr{F}_{\omega,D}(\omega_\phi) &= -\frac{d^2}{dt^2}F^0_{\omega,D}(\phi)=\frac{\int_D(\omega+\partial\bar{\partial}\Phi)^n/n!}{dt\wedge d\bar{t}}\\ &= \int_D \ddot{\phi}\omega_{KE}^{n-1}/(n-1)!-\int_D\partial\dot{\phi}\wedge\bar{\partial}\dot{\phi}\wedge\omega_{KE}^{n-2}/(n-2)!\\ &= \int_D(|\nabla\dot{\phi}|^2_{\omega_{KE}}-|\nabla^D\dot{\phi}|^2_{\omega_{KE}|_D})\omega_{KE}^{n-1}/(n-1)!\\ &= \int_X|(\nabla\dot{\phi})^\perp|^2_{\omega_{KE}}\omega_{KE}^{n-1}/(n-1)!\geq 0. \qquad (37)\end{aligned}$$

□

Lemma 7.2. *We have the following different formulas for the Hessian of $\mathscr{F}_{\omega,D}$ on $\mathscr{M}_{KE}$.*

$$\begin{aligned}Hess\mathscr{F}(\theta,\theta) &= \int_D|\nabla\theta^\perp|^2\omega_{KE}^{n-1}/(n-1)! \qquad (38)\\ &= \lambda\int_X\theta^2\omega_{KE}^n/n!+\int_X(-\theta^2+\theta^i\theta_i)(\lambda\phi-\log\|s\|_h^2)\omega_{KE}^n/n! \qquad (39)\\ &= \lambda\int_X\theta^2\omega_{KE}^n/n!+\int_X(\theta^2-\theta^i\theta_i)(\log\|s\|^2_{he^{-\lambda\phi}})\omega_{KE}^n/n!. \qquad (40)\end{aligned}$$

Proof. The first identity follows from (37) because $\theta=\frac{\partial\phi}{\partial t}|_{t=0}$. For the second identity, we shall give two calculations. For $\theta\in\Lambda_1$, $\nabla\theta$ is a holomorphic vector field generating a one parameter subgroup σ_t in $Aut(X)$. Let $\sigma_t^*\omega_{KE}=\omega_{KE}+\sqrt{-1}\partial\bar{\partial}\phi_t$. Then ϕ_t satisfies the geodesic equation: $\ddot{\phi}-|\nabla\dot{\phi}|^2_{\omega_\phi}=0$ with initial velocity $\frac{d}{dt}\phi|_{t=0}=\theta$.

- (calculation 1)

$$\begin{aligned}
\frac{d^2}{dt^2}\mathscr{F}_{\omega,D} &= \frac{d^2}{dt^2}F^0_{\omega,D}(\phi) = \int_D \ddot{\phi}\omega_{KE}^{n-1}/(n-1)! - \int_D \partial\dot{\phi}\wedge\bar{\partial}\dot{\phi}\wedge\omega_{KE}^{n-2}/(n-2)! \\
&= \int_X (\sqrt{-1}\partial\bar{\partial}\log\|s\|_h^2 + \lambda(-\sqrt{-1}\partial\bar{\partial}\phi + \omega_{KE}))\wedge(\ddot{\phi}\omega_{KE} - (n-1)\partial\dot{\phi}\wedge\bar{\partial}\dot{\phi})\wedge\omega_{KE}^{n-2}/(n-1)! \\
&= \int_X (\log\|s\|^2 - \lambda\phi)(\sqrt{-1}\partial\bar{\partial}\ddot{\phi}\wedge\omega_{KE}^{n-1} + (n-1)\partial\bar{\partial}\dot{\phi}\wedge\partial\bar{\partial}\dot{\phi}\wedge\omega_{KE}^{n-2})/(n-1)! + \\
&\quad +\lambda\int_X (\ddot{\phi}\omega_{KE}^n - (n-1)\partial\dot{\phi}\wedge\bar{\partial}\dot{\phi}\wedge\omega_{KE}^{n-1})/(n-1)! \\
&= \int_X (\log\|s\|^2 - \lambda\phi)(\Delta|\nabla\theta|^2 + (\Delta\theta)^2 - \theta_{i\bar{j}}\theta^{i\bar{j}})\omega_{KE}^n/n! + \lambda\int_X \theta^i\theta_i\omega_{KE}^n/n! + \\
&\quad +\lambda\int_X (\ddot{\phi} - |\nabla\dot{\phi}|^2_{\omega_{KE}})\omega_{KE}^n/(n-1)! \\
&= \int_X (\log\|s\|^2_{he^{-\lambda\phi}})(\theta^2 - \theta^i\theta_i) + \lambda\int_X \theta^2\omega_{KE}^n/n!.
\end{aligned}$$

- (Calculation 2)

$$\begin{aligned}
\left.\frac{d^2}{dt^2}\right|_{t=0}\omega_t^n &= \frac{d}{dt}(\Delta\dot{\phi}\omega_t^n) = (\dot{\Delta}\dot{\phi} + \Delta\ddot{\phi} + (\Delta\dot{\phi})^2)\omega_{KE}^n \\
&= (-\theta^{i\bar{j}}\theta_{i\bar{j}} + (\theta^i\theta_i)_j{}^j + \theta^2)\omega_{KE}^n \\
&= (-\theta^{i\bar{j}}\theta_{i\bar{j}} + \theta^i{}_j\theta_i{}^j + \theta^i{}_j{}^j\theta_i + \theta^2)\omega_{KE}^n/n! \\
&= (\theta^2 - \theta^i\theta_i)\omega_{KE}^n
\end{aligned}$$

$$\begin{aligned}
Hess\mathscr{F}(\theta,\theta) &= \frac{d^2}{dt^2}\left(\lambda(I-J)_\omega(\omega_t) + \int_X \log\|s\|_h^2\omega_t^n\right) \\
&= -\lambda\int_X \dot{\phi}\Delta\dot{\phi}\omega_{KE}^n/n! + \int_X (-\lambda\phi + \log\|s\|_h^2)\frac{d^2}{dt^2}\omega_t^n/n! \\
&= \lambda\int_X \theta^2\omega_{KE}^n/n! + \int_X (\log\|s\|^2_{he^{-\lambda\phi}})(\theta^2 - \theta^i\theta_i)\omega_{KE}^n/n!.
\end{aligned}$$

□

Lemma 7.3. *Under the assumption that $\lambda \geq 1$, $Hess\mathscr{F}$ is non-degenerate at any point $\omega_{KE} \in \mathscr{M}_{KE}$.*

Proof. We have seen $Hess\mathscr{F}$ is non-negative at any point $\omega_{KE} \in \mathscr{M}_{KE}$ using formula (38). $Hess\mathscr{F}$ is degenerate if and only if $\int_D |(\nabla\theta)^\perp|^2\omega_{KE}^{n-1}/(n-1)! = 0$. This happens if and only if $(\nabla\theta)^\perp \equiv 0$ on D, i.e. when $\nabla\theta$ is tangent to D. However, since we assume $\lambda \geq 1$, we are in the log-canonically-polarized or log-Calabi-Yau case, there can not be holomorphic vector fields tangent to D by Corollary 4.3 (see also [51]). So the lemma follows. □

Lemma 7.4. *When restricted to $\mathscr{M}_{KE}$, there exists a unique minimum ω_{KE}^D of $\mathscr{F}_{\omega,D}(\omega_\phi)$.*

Proof. By the previous Lemma, $\mathscr{F}_{\omega,D}$ is a convex functional on the space $\mathscr{M}_{KE} \cong G^{\mathbb{C}}/G$. To prove the existence of critical point, we only need to show it's proper on $G^{\mathbb{C}}/G$. Because we assumed $\lambda \geq 1$ and there exists Kähler-Einstein on X, by Theorem 4.1, $\mathcal{M}_{\omega,(1-\beta)D}$ is proper for $\beta \in (0,1)$. Because the Mabuchi energy is constant on $\mathscr{M}_{KE}$, by equality (36), $\mathscr{M}_{\omega,(1-\beta)D} = (1-\beta)\mathscr{F}_{\omega,D}$+constant is proper on $\mathscr{M}_{KE}$. □

Write $\omega_{KE}^D = \omega + \sqrt{-1}\partial\bar{\partial}\phi_{KE}^D$, then it satisfies the critical point equation

$$\int_X (\log\|s\|^2 - \lambda\phi_{KE}^D)\psi(\omega_{KE}^D)^n = 0.$$

for any $T_{\omega^D_{KE}}\mathscr{M}_{KE} \cong \Lambda_1(\omega^D_{KE})/\mathbb{C}$. In other words, $\lambda\phi^D_{KE} - \log\|s\|^2 \in \Lambda_1^{\perp}$.

Proposition 7.1. *As $\beta \to 1$, the conical Kähler-Einstein metrics ω_β converges to a unique smooth Kähler-Einstein metric $\omega^D_{KE} \in \mathscr{M}_{KE}$.*

Proof. Recall that the conical Kähler-Einstein equation is

$$(\omega + \sqrt{-1}\partial\bar{\partial}\phi)^n = e^{h_\omega - r(\beta)\phi}\frac{\omega^n}{\|s\|^{2(1-\beta)}}.$$

Any $\omega_{KE} = \omega + \sqrt{-1}\partial\bar{\partial}\phi_{KE} \in \mathscr{M}_{KE}$ satisfies the equation

$$(\omega + \sqrt{-1}\partial\bar{\partial}\phi_{KE})^n = e^{h_\omega - \phi_{KE}}\omega^n.$$

By Lemma 7.4, there exists a unique minimum ω^D_{KE} of the functional $\mathscr{F}_{\omega,D}$ on $\mathscr{M}_{KE}$. We will choose $\omega_{KE} = \omega^D_{KE}$ in the following argument. Divide the above two equations to get

$$\log\frac{(\omega + \sqrt{-1}\partial\bar{\partial}\phi)^n}{(\omega + \sqrt{-1}\partial\bar{\partial}\phi_{KE})^n} = \phi_{KE} - r(\beta)\phi - (1-\beta)\log\|s\|^2_h.$$

Let $\phi = \phi_{KE} + \psi$ and $\psi = \theta + \psi'$ with $\theta \in \Lambda_1$ and $\psi' \in \Lambda_1^{\perp}$, then

$$\log\frac{(\omega_{KE} + \sqrt{-1}\partial\bar{\partial}(\theta + \psi'))^n}{\omega^n_{KE}} + r(\beta)(\theta + \psi') = (1-\beta)(\lambda\phi_{KE} - \log\|s\|^2_h). \tag{41}$$

We use Bando-Mabuchi's bifurcation method to solve the equation for β close to 1. First project to Λ_1 to get

$$(1 - P_0)\left(\log\frac{(\omega_{KE} + \sqrt{-1}\partial\bar{\partial}(\theta + \psi'))^n}{\omega^n_{KE}}\right) + r(\beta)\psi' = (1-\beta)(\lambda\phi_{KE} - \log\|s\|^2_h). \tag{42}$$

The equation is satisfied for $(\beta, \psi, \theta) = (1, 0, 0)$. The linearization of the left side of this equation with respect to ψ' is

$$(1 - P_0)(\tilde{\Delta}_\theta + r(\beta))\psi'$$

where $\tilde{\Delta}_\theta$ is the Laplacian with respect to $\omega_{KE} + \sqrt{-1}\partial\bar{\partial}\theta$. Since $\Lambda_1 = Ker(\Delta_0 + 1)$, there exists a positive constant $\delta > 0$, such that

$$(1 - P_0)(-\Delta_{\omega_{KE}} - 1) \geq \delta > 0.$$

By continuity, it's easy to see that

$$(1 - P_0)(-\tilde{\Delta}_\theta - r(\beta)) \geq \delta/2 > 0.$$

for (β, θ) close to $(1, 0)$. In other words, the inverse of $(1 - P_0)(\tilde{\Delta}_\theta + r(\beta))$ has uniformly bounded operator norm for (β, θ) close to $(1, 0)$. So by implicit function theorem, there exists solution $\psi'_{\beta,\theta}$ for β near 1 and θ small. Now to solve the equation (41), we only need to solve the following equation

$$P_0\left(\log\frac{(\omega_{KE} + \sqrt{-1}\partial\bar{\partial}(\theta + \psi'))^n}{\omega^n_{KE}}\right) = -r(\beta)\theta. \tag{43}$$

Let $\sigma^*\omega_{KE} = \omega_{KE} + \sqrt{-1}\partial\bar{\partial}(\theta + \psi'_{1,\theta})$. We have the identity

$$\log\frac{(\omega_{KE} + \sqrt{-1}\partial\bar{\partial}(\theta + \psi'_{1,\theta}))^n}{\omega^n_{KE}} = -(\theta + \psi'_{1,\theta})$$

Let $\psi'_{\beta,\theta} = \psi'_{1,\theta} + (1-\beta)\xi_{\beta,\theta}$. We can rewrite the equation (41) in the following form

$$\begin{aligned}
&\log\frac{(\omega_{KE} + \sqrt{-1}\partial\bar{\partial}(\theta + \psi'_{1,\theta} + (1-\beta)\xi_{\beta,\theta}))^n}{\omega^n_{KE}} = \\
= \;& -(1 - \lambda(1-\beta))(\theta + \psi'_{1,\theta} + (1-\beta)\xi_{\beta,\theta}) + (1-\beta)(\lambda\phi_{KE} - \log\|s\|^2_h) \\
= \;& \log\left(\frac{\omega^n_\theta}{\omega^n_{KE}}\right) + (1-\beta)\lambda(\theta + \psi'_{1,\theta}) - (1-\beta)r(\beta)\xi_{\beta,\theta} + (1-\beta)(\lambda\phi_{KE} - \log\|s\|^2_h).
\end{aligned}$$

where $\omega_\theta = \omega_{KE} + \sqrt{-1}\partial\bar\partial(\theta + \psi'_{1,\theta})$. In particular, it's easy to see that (42) is equivalent to

$$\frac{1}{1-\beta}(1-P_0)\left(\log\frac{(\omega_{KE}+\sqrt{-1}\partial\bar\partial(\theta+\psi'_{1,\theta}+(1-\beta)\xi_{\beta,\theta}))^n}{(\omega_{KE}+\sqrt{-1}\partial\bar\partial(\theta+\psi'_{1,\theta}))^n}\right) = \lambda\psi'_{1,\theta} - r(\beta)\xi_{\beta,\theta} + (\lambda\phi_{KE} - \log\|s\|_h^2).$$

Let $\beta \to 1$ to get

$$(1-P_0)\left((\Delta_\theta+1)\xi_{1,\theta}\right) - \lambda\psi'_{1,\theta} = \lambda\phi_{KE} - \log\|s\|_h^2.$$

where Δ_θ is the Laplacian with respect to the metric ω_θ. Since $Im(\Delta_0+1) = Ker(\Delta_0+1)^\perp = \Lambda_1^\perp$, so in particular,

$$(\Delta_0+1)\xi_{1,0} = \lambda\phi_{KE} - \log\|s\|_h^2. \tag{44}$$

Now the equation (43) is equivalent to

$$P_0\left(\frac{1}{1-\beta}\log\frac{(\omega_{KE}+\sqrt{-1}\partial\bar\partial(\theta+\psi'_{1,\theta}+(1-\beta)\xi_{\beta,\theta}))^n}{(\omega_{KE}+\sqrt{-1}\partial\bar\partial(\theta+\psi'_{1,\theta}))^n}\right) - \lambda\theta = 0 \tag{45}$$

Denote by $\Gamma(\beta,\theta)$ the term on the left side, Then

$$\Gamma(1,0) = 0, \quad \Gamma(1,\theta) = P_0(\Delta_\theta\xi_{1,\theta}) - \lambda\theta.$$

Let $\theta(t) = t\theta \in \Lambda_1$, for any $\theta' \in \Lambda_1 = Ker(\Delta_0+1)$,

$$\begin{aligned}\int_X \frac{d}{dt}\Gamma(1,\theta)\Big|_{t=0}\theta'\omega_{KE}^n/n! &= -\lambda\int_X \theta\theta'\omega_{KE}^n/n! + \int_X(\dot\Delta_\theta\xi_{1,0} + \Delta_0\dot\xi_{1,0})\theta'\omega_{KE}^n/n!\\ &= -\lambda\int_X \theta\theta'\omega_{KE}^n/n! + \int_X -\theta^{i\bar j}(\xi_{1,0})_{i\bar j}\theta'\omega_{KE}^n/n!.\end{aligned}$$

Let $\xi = \xi_{1,0} \in \Lambda_1^\perp$. As the calculation in [4], we have

$$\begin{aligned}\int_X \theta^{i\bar j}\xi_{i\bar j}\theta'\omega_{KE}^n/n! &= -\int_X(\theta^i\xi_{i\bar j}{}^{\bar j}\theta' + \theta^i\xi_{i\bar j}\theta'^{\bar j})\omega_{KE}^n/n! = -\int_X(\theta^i\xi_{\bar j}{}^{\bar j}{}_i\xi' - \theta^i\xi_i\theta'^{\bar j}_{\bar j})\omega_{KE}^n/n!\\ &= -\int_X \theta^i((\Delta+1)\xi)_i\theta'\omega_{KE}^n/n! = \int_X(-\theta\theta' + \theta^i\theta'_i)(\Delta+1)\xi\omega_{KE}^n/n!\\ &= -\int_X(\theta\theta' - \theta^i\theta'_i)(\lambda\phi - \log\|s\|_h^2)\omega_{KE}^n/n!\end{aligned}$$

So

$$\begin{aligned}D_2\Gamma(1,0)(\theta)\theta' &= -\lambda\int_X \theta\theta'\omega_{KE}^n/n! + \int_X(\theta\theta' - \theta^i\theta'_i)(\lambda\phi - \log\|s\|_h^2)\omega_{KE}^n/n!\\ &= -Hess\mathscr{F}(\theta,\theta'). \quad (\text{ by equation (40)})\end{aligned}$$

By Lemma 7.3, $D_2\Gamma(1,0)$ is invertible, so by implicit function theorem, (45) is solvable for β close to 1. So we get conical Kähler-Einstein metrics ω_β for β close to 1 and by continuity ϕ_β converges to ϕ_{KE}^D as $\beta \to 1$. ☐

Remark 7.1. *As in [4], we can continue to calculate:*

$$\begin{aligned}\int_X \theta^{i\bar{j}}\xi_{i\bar{j}}\theta'\omega_{KE}^n/n! &= -\int_X(\theta\theta'-\theta^i\theta'_i)(\lambda\phi-\log\|s\|_h^2)\omega_{KE}^n/n!\\ &= \frac{1}{2}\int_X(\Delta\theta\theta'+\theta\Delta\theta'+\theta^i\theta'_i+\theta_i\theta'^i)(\Delta+1)\xi\omega_{KE}^n/n!\\ &= \frac{1}{2}\int_X(\Delta(\theta\theta'))(\Delta+1)\xi\omega_{KE}^n/n! = \frac{1}{2}\int_X\theta\theta'\Delta(\Delta+1)\xi\omega_{KE}^n/n!\\ &= \frac{1}{2}\int_X\theta\theta'\Delta(\lambda\phi_1-\log\|s\|_h^2)\omega_{KE}^n/n!\\ &= \frac{1}{2}\int_X\theta\theta' n\sqrt{-1}\partial\bar{\partial}(\lambda\phi_1-\log\|s\|_h^2)\wedge\omega_{KE}^{n-1}/n!\\ &= \frac{1}{2}n\lambda\int_X\theta\theta'\omega_{KE}^n/n!-\frac{1}{2}\int_D\theta\theta'\omega_{KE}^{n-1}/(n-1)!.\end{aligned}$$

so that

$$D_2\Gamma(1,0)(\theta)\theta' = -\lambda(1+n/2)\int_X\theta\theta'\omega_{KE}^n/n!+\frac{1}{2}\int_D\theta\theta'\omega_{KE}^{n-1}/(n-1)!.$$

However, it seems not straightforward to see that $D_2\Gamma(1,0)$ is nondegenerate using this formula.

Remark 7.2. *One reason why we packed all the conical spaces together in the space of admissible functions is because that we need to work in different function space corresponding to different cone angles. Strictly speaking, there are subtleties in applying implicit functional theorem in this setting. However, we expect one can generalize Donaldson's argument to validate the application of implicit function theorem.*

8. RELATIONS TO SONG-WANG'S WORK

In this section, we will briefly explain Song-Wang's results and derive one of its implications. For one thing, they also observe the interpolation property for the log-Ding-energy. Secondly, their prominent idea of considering pluri-anticanonical sections corresponds to the $\lambda \geq 1$ case in our paper. Recall that $R(X)$ in the introduction (see (2)) is defined to be the greatest lower bound of Ricci curvature of smooth Kähler metrics in $c_1(X)$. $R(X)$ was studied in ([56],[54],[33],[36],[51]). Donaldson ([20]) made the following conjecture

Conjecture 8.1 (Donaldson). *Let $D \in |-K_X|$ be a smooth divisor, then there exists a conical Kähler-Einstein metric on $(X,(1-\beta)D)$ if and only if $\beta \in (0, R(X))$.*

Song-Wang proved a weak version of Donaldson's conjecture by allowing pluri-anticanonical divisor and its dependence on β. Translating their result in our notations, they proved

Theorem 8.1 (Song-Wang,[51]). *For any $\gamma \in (0, R(X))$ there exists a large $\lambda \in \mathbb{Z}$ and a smooth divisor $D \in |\lambda K_X^{-1}|$ such that there exists a conical Kähler-Einstein metric on $(X, \lambda^{-1}(1-\gamma)D)$.*

Remark 8.1. *Note that in general, λ and D may depend on γ. γ is related to the cone angle parameter β by the relation $\lambda^{-1}(1-\gamma) = 1-\beta$ or equivalently, $\gamma = r(\beta) = 1-\lambda(1-\beta)$.*

The proof of this theorem can be explained through the Hölder's inequality

$$\int_X e^{h_\omega-\gamma\phi}\frac{\omega^n}{n!\|s\|^{2(1-\gamma)/\lambda}} \leq \left(\int_X e^{p(h_\omega-\gamma\phi)}\omega^n/n!\right)^{1/p}\left(\int_X \|s\|^{-2q(1-\gamma)/\lambda}\omega^n/n!\right)^{1/q}$$

where $p^{-1}+q^{-1}=1$. To make contact with the invariant $R(X)$, one choose $p=\frac{t}{\gamma}$ for any $t \in (\gamma, R(X))$. (This is related to the characterization of $R(X)$ through the properness of twisted Ding-energy as in [36]) Then $q=(1-p^{-1})^{-1}=\frac{t}{t-\beta}$. Now the integrability of the second integral on the right gives the restriction on λ: $\frac{2q(1-\gamma)}{\lambda}-1<1$. This gives the the lower bound of λ in Song-Wang's theorem.

$$\lambda>(1-\gamma)\frac{R(X)}{R(X)-\gamma}.$$

One other important result Song-Wang proved is the construction of toric conical Kähler-Einstein metrics can be combined with the strategy in our paper to prove a version of Donaldson's conjecture on toric Fano manifolds. We will explain this briefly.

Any toric Fano manifold $X_\triangle$ is determined by a reflexive lattice polytope $\triangle\subset\mathbb{R}^n$ containing only O as the interior lattice point. For any $P\in\mathbb{R}^n$, P determines a toric $\mathbb{R}$-divisor $D_P\sim_{\mathbb{R}}-K_X$. More concretly, assume that the polytope is defined by the inequalities $l_j(x)=\langle x,\nu_j\rangle+a_j\geq 0$. Then $D_P=\sum_j l_j(P)D_j$. If $P\in\overline{\triangle}$ is a rational point, then for any integer λ such that λP is an integral lattice point, there exist a genuine holomorphic section s_P^λ of $-\lambda K_X$ and an integral divisor λD_P.

Let P_c be the barycenter of $\triangle$, then the ray $\overrightarrow{P_cO}$ intersect the boundary $\partial\triangle$ at a unique point Q. Note that in general, Q is a rational point. In [33], the first author proved $R(X)$ is given by

$$R(X)=\frac{|\overline{OQ}|}{|\overline{P_cQ}|}. \tag{46}$$

For any $\gamma\in[0,1]$, define $P_\gamma=-\frac{\gamma}{1-\gamma}P_c$. Then $P_\gamma\in\overline{\triangle}$ if and only if $\gamma\in[0,R(X)]$, which is also equivalent to D_{P_γ} being effective. In particular, $P_{R(X)}=Q$. Using these notations, Song-Wang proved the following theorem by adapting the method in Wang-Zhu's work ([64]) on the existence of Kähler-Ricci solitons on toric Fano manifolds.

Theorem 8.2 (Song-Wang,[51])**.** *For any $\gamma\in[0,1]$, there exists toric solution to the following equation:*

$$Ric(\omega)=\gamma\omega+(1-\gamma)\{D_{P_\gamma}\}.$$

When $\gamma\in[0,R(X)]$ is rational, then the solution ω_γ is a conical Kähler-Einstein metric on $(X,(1-\gamma)D_{P_\gamma})$. In particular, when $\gamma=R(X)$, there exists a conical Kähler-Einstein metric on $(X,(1-R(X))D_Q)$.

Example 8.1. *The above theorem can be generalized to toric orbifold case. (See [50] for related work where the Kähler-Ricci soliton on toric Fano orbifolds was considered) We will illustrate this by showing the conical Kähler-Einstein on $X=\mathbb{P}(1,1,4)$ considered in section 6 in the toric language. The polytope determining $(X,-K_X)$ is the following rational polytope $\triangle$. Note that $-2K_X$ is Cartier because $2\triangle$ is a lattice polytope. $Q=(-1,1/2)$, $P_c=(1,-1/2)$. So $R(X)=|\overline{OQ}|/|\overline{P_cQ}|=1/2$. $D_Q=3/2D$, where the divisor D corresponds to the facet $\overline{AB}$. The conical Kähler-Einstein satisfies the equation:*



$$Ric(\omega)=\frac{1}{2}\omega+(1-\frac{1}{2})\cdot\frac{3}{2}D.$$

So the cone angle along D is $2\pi\beta$ with $\beta=1-3/4=1/4$.

Now we show that Song-Wang's nice existence result implies Theorem 1.2.

Proof of Theorem 1.2. Let $\mathcal{F}_Q$ be the minimal face of $\triangle$ containing Q. For any $\lambda \in \mathbb{Z}$ such that λQ is an integral point, define a set of rational points by

$$\mathcal{R}(Q,\lambda) = \{Q\} \bigcup \left((\overline{\triangle} \setminus \overline{\mathcal{F}_Q}) \bigcap \frac{1}{\lambda}\mathbb{Z}^n \right).$$

Then we define the linear system $\mathscr{L}_\lambda$ to be the linear subspace spanned by the holomorphic sections corresponding to rational points in $\mathcal{R}(Q,\lambda)$:

$$\mathscr{L}_\lambda = Span_{\mathbb{C}} \left\{ s_P^\lambda; P \in \mathcal{R}(Q,\lambda) \right\}.$$

Choose any general element $D \in \mathscr{L}_\lambda$, the coefficient of the term s_Q^λ is nonzero. Because Q is a vertex of the convex hull of $\mathcal{R}(Q,\lambda)$, there exists a $\mathbb{C}^*$ action denoted by $\sigma(t)$ contained in the torus action, such that

$$\lim_{t\to 0} \sigma(t)^* D = \lambda D_Q.$$

In this way, we construct a degeneration $(\mathcal{X}, \frac{1-R(X)}{\lambda}\mathcal{D}, K_{\mathcal{X}}^{-1})$ with $\mathcal{X} = X \times \mathbb{C}$ and $\mathcal{D}_t = \sigma_t^* D$. By Song-Wang's theorem in [51], the central fibre $(\mathcal{X}_0, \frac{1-R(X)}{\lambda}(\lambda D_Q)) = (X, (1-R(X))D_Q)$ has conical Kähler-Einstein metric. So we use Theorem 1.4 to get the lower bound of $\mathcal{M}_{X, \frac{1-R(X)}{\lambda}D}$. (As explained in the proof of Theorem 1.4, in the present case, since $\mathcal{X} = X \times \mathbb{C}$, we just need to use the trivial geodesic and apply Berndtsson's result in [10] to get the subharmonicity and complete the proof) On the other hand, because $\lambda \geq 1$, we can use the interpolation result in Proposition 1.1 to see that $\mathcal{M}_{X,\frac{1-\gamma}{\lambda}D}$ is proper for any $\gamma \in (0, R(X))$ (actually for any $\gamma \in (1-\lambda, R(X))$. So there exists a conical Käler-Einstein metric on $(X, \frac{1-\gamma}{\lambda}D)$ for any $\gamma \in (0, R(X))$. There can not be conical Kähler-Einstein metric for $\gamma \in (R(X), 1)$ is easy to get because the twisted energy is bounded from below by the log-Ding-energy. For details, see [51] and also [36]. The non-existence for $\gamma = R(X)$ is implied by Donaldson's openness theorem in [20] (see Theorem 2.1), since otherwise there exists conical Kähler-Einstein for some $\gamma \in (R(X), 1)$. ☐

Remark 8.2. *The smoothness of the generic member seems to be more subtle than we first thought. We will discuss this a little bit using standard toric geometry. For this, we first denote* $\{H_i\}_{i=1}^N$ *to be the set of codimensional 1 face (i.e. facet) of* $\triangle$. *Define*

$$\mathcal{B}(\mathcal{F}_Q) = \left(\bigcup_{\mathcal{F}_Q \not\subset H_i} H_i \right) \bigcap \mathcal{F}_Q.$$

Now it's easy to see that the base locus of $\mathscr{L}_\lambda$ *is equal to*

$$\mathbb{B}_Q = \bigcup_{\sigma \subset \mathcal{B}(\mathcal{F}_Q)} X_\sigma.$$

Here for any face of $\triangle$ *we denote* X_σ *to be the toric subvariety determined by* σ. *Indeed, this follows from the following fact: if* P *is any lattice point and* $\mathcal{F}_P$ *is the minimal face containing* P. *Define*

$$\boldsymbol{Star}(\mathcal{F}_P) = \bigcup_{\mathcal{F}_P \subset \sigma} \sigma.$$

where σ *ranges over all the (closed) faces of* $\triangle$. *(including* $\triangle$ *itself). Then the zero set of the corresponding holomorphic section* s_P *is the toric divisor corresponding to the set*

$$\overline{\triangle} \setminus (\boldsymbol{Star}(\mathcal{F}_P))^\circ = \bigcup_{\mathcal{F}_P \not\subset H_i} H_i \subset \partial\triangle.$$

By Bertini's Theorem ([26]), the generic element $D \in \mathscr{L}_\lambda$ *is smooth away from* $\mathbb{B}_Q$. *To analyze the situation near* $\mathbb{B}_Q$, *fix any vertex* P *of* $\mathcal{F}$. *We can choose integral affine coordinates* $\{x_i\}_{i=1}^n$

such that

$$\mathcal{F}_Q = \bigcap_{i=m+1}^{n} \{x_i = 0\}.$$

We can also write $Q = \lambda(d_1, \dots, d_m, 0, \dots, 0)$ *with* λd_i *being positive integers. On the other hand, by standard toric geometry, the normal fan of* $\triangle$ *at* P *determines an affine chart* $\mathcal{U}_P$ *on* X*. There exists complex coordinate* $\{z_i\}_{i=1}^n$ *such that* $X_{\mathcal{F}_Q} \cap \mathcal{U}_P = \{z_{m+1} = 0, \dots, z_n = 0\}$*. Locally, the generic member* D *in* $\mathscr{L}_\lambda$ *is given by the equation of the form:*

$$\prod_{i=1}^{m} a_i z_i^{\lambda d_i} + \sum_{j=m+1}^{n} b_j z_j (1 + f_j(z_1, \dots, z_m)) + \sum_{j,k=m+1}^{n} c_{jk} z_j z_k g(z_1, \dots, z_n)).$$

where $a_i, b_j \neq 0$*. If we delete the lattice points corresponding to terms* $z_j f_j(z_1, \dots, z_m)$*, then* C *would be smooth near* $X_{\mathcal{F}_Q} \cap \mathcal{U}_P$*. Since* $\mathbb{B}_Q \subset X_{\mathcal{F}_Q}$ *and* $\mathcal{U}_P$ *covers* $X_{\mathcal{F}_Q}$ *as* P *ranges over all the vertices of* $\mathcal{F}_Q$ *we conclude that* D *is smooth at points in* $\mathbb{B}_Q$ *as well. This certainly puts a lot of restriction on the sub-linear system. However, even if we don't delete these lattice points, the generic member in* $\mathscr{L}_\lambda$ *could be smooth. For example, this is the case when* $\mathcal{F}_Q$ *has dimension* ≤ 1 *in which case the base locus consists of isolated points. In particular, this is true when the toric variety has dimension* ≤ 2.

Remark 8.3. *The degeneration behavior in the toric case is closely related to the study of degenerations in* [34] *where the current* D_{P_γ} *is replaced by* $(1-\gamma)\omega$ *with* ω *being a smooth reference metric.*

Example 8.2. *Let* $X = Bl_p\mathbb{P}^2$*. Let* $[Z_0, Z_1, Z_2]$ *be homogeneous coordinate on* $\mathbb{P}^2$*. We can assume* $p = (1, 0, 0) \in \mathbb{C}^2 = \{Z_0 \neq 0\} \subset \mathbb{P}^2$*. Let* $\pi : X \to \mathbb{P}^2$ *be the blow down of exceptional divisor* E*. For simplicity we use* H *to denote both the hyperplane class on* $\mathbb{P}^2$ *and its pull-back on* X*. Then* $-K_X = 3H - E$ *and* $-2K_X = 6H - 2E$*. So divisors in* $|-2K_X|$ *correspond to the sextic curves on* $\mathbb{P}^2$ *whose vanishing order at* 0 *is at least 2. More precisely, if* C *is such a curve representing* $6H$*, then the corresponding divisor* $D(C)$ *in* $|-2K_X| = |6H - 2E|$ *is the strict transform of* C*. In toric language* X *is determined by the following polytope:*





The invariant $R(X) = 6/7$ *was calculated in* [54] *and* [33]*. Since the point* $Q = (-1/2, -1/2)$*, it's easy to see that* $D_Q = 1/2(F_1 + F_2) + 2D_\infty$*. By Song-Wang* [51]*, there is a conical Kähler-Einstein metric on* $(X, (1 - R(X))D_Q) = (X, 1/7D_Q)$*.*

Now λQ *is integral when* λ *is even. The generic divisors in the linear system* $\mathscr{L}_2$ *correspond to the sextic curves given by degree* 6 *homogeneous polynomial of the form*

$$C : Z_0^4 Z_1 Z_2 + \sum_{i=0}^{3} \sum_{j=0}^{6-i} a_{ij} Z_0^i Z_1^j Z_2^{6-i-j} = 0.$$

Let σ_t *be the* $\mathbb{C}^*$*-action given by*

$$(Z_0, Z_1, Z_2) \to (Z_0, t^{-1}Z_1, t^{-1}Z_2).$$

Then $\lim_{t\to 0} \sigma_t \cdot C = \{Z_0^4 Z_1 Z_2 = 0\}$*. Equivalently, by taking strict transform, we get* $\lim_{t\to 0} \sigma_t \cdot D(C) = 2D_Q$*. The same argument applies to* $\lambda = 2m$ *being even, where the divisors in* $\mathscr{L}_{2m}$

correspond to the degree $6m$ *curves of the form:*

$$Z_0^{4m}Z_1^mZ_2^m+\sum_{i=0}^{4m-1}\sum_{j=0}^{6m-i}a_{ij}Z_0^iZ_1^jZ_2^{6m-i-j}=0.$$

Note that the strict transform of such generic curves are smooth at the base locus $\mathbb{B}_Q=B_1\cup B_2$ *and so smooth everywhere.*

Remark 8.4. *From the above discussion, we see that when* λ *is even, the divisor degenerates while the ambient space stays the same. The case when* $\lambda=1$*, or more generally when* λ *is odd, is still open. From the point of view in our strategy, the problem is that the right degeneration to conical Kähler-Einstein pair is still missing. In this case, we expect the degeneration also happens to the ambient space, similar with the degree 2 plane curve case studied in section 6.*

References

[1] Claudio Arezzo; Alessandro Ghigi; Gian Pietro Pirola, *Symmetries, quotients and Kähler-Einstein metrics.* J. Reine Angew. Math. 591 (2006), 177200.

[2] Thierry Aubin: Equations du type Monge-Ampère sur les variétés compactes, C.R.Acad. Sc. Paris, 283, 1976, pp. 119-121.

[3] Shigetoshi Bando: The K-energy map, almost Einstein Kähler metrics and an inequality of the Miyaoka-Yau type. Tohuku Mathematical Journal 39 (1987): 231-235.

[4] Shigetoshi Bando, Toshiki Mabuchi: Uniqueness of Einstein Kähler metrics modulo connected group actions, *Algebraic geometry*, Sendai 1985, Advanced Studies in Pure Mathematics 10, 1987: 11-40

[5] Eric Bedford, and B.A.Taylor: The Dirichlet Problem for a complex Monge-Ampère equation, Inventiones Math. 37, 1-44(1976).

[6] Robert J. Berman: A thermodynamic formalism for Monge-Ampère equations, Moser-Trudinger inequalities and Kähler-Einstein metrics arXiv:1011.3976

[7] Robert J. Berman: K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics, arXiv:1205.6214.

[8] Robert J. Berman, Sébastien Boucksom, Philippe Eyssidieux, Vincent Guedj, Ahmed Zeriahi, Kähler-Ricci flow and Ricci iteration on log-Fano varieties, arXiv:1111.7158.

[9] Bo Berndtsson: Subharmonicity properties of the Bergman kernel and some other functions associated to pseudoconvex domains, arXiv:0505469, to appear in Ann. Inst. Fourier.

[10] Bo Berndtsson: A Brunn-Minkowski type inequality for Fano manifolds and the Bando-Mabuchi uniqueness theorem,

[11] Bo Berndtsson, and Mihai Paun: Bergman kernels and the pseudoeffectivity of relative canonical bundles. Duke Math. J. Volume 145, Number 2 (2008), 341-378.

[12] Zbigniew Błocki: On geodesics in the space of Kähler metric.

[13] Simon Brendle: Ricci flat Kähler metics with edge singularities, arXiv:1103.5454.

[14] L.Caffarelli, J.J.Kohn, L.Nirenberg, J.Spruck, The Dirichlet problem for non-linear second order elliptic equations II: Complex Monge-Ampère, and uniformly elliptic equations. Comm. Pure Appl. Math. 38 (1985), 209-252.

[15] Xiuxiong Chen, Space of Kähler metric, J. Differential Geometry, 56 (2000) 189-234.

[16] Xiuxiong Chen, Space of Kähler metrics (IV)–On the lower bound of the K-energy, arXiv:0809.4081.

[17] Diego Conti: Cohomogeneity one Einstein-Sasaki 5-manifolds. Comm. Math. Phys. 274 (2007), no. 3, 751-774.

[18] Wei-Yue Ding: Remarks on the existence problem of positive Kähler-Einstein metrics. Math. Ann. 282, 463-471 (1988)

[19] S.K. Donaldson: Scalar curvature and stability of toric varieties, Jour. Differential Geometry 62 289-349 (2002)

[20] S.K. Donaldson: Kähler metrics with cone singularities along a divisor. arXiv:1102.1196

[21] S.K. Donaldson, and Song Sun: Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry. arXiv:1206.2609.

[22] P. Eyssidieux, V. Guedj, and A. Zeriahi: Singular Kähler-Einstein metrics, Journal of American Mathematical Society, Vol 22, No. 3, 2009, 607-639.

[23] Akito Futaki: An obstruction to the existence of Einstein Kähler metrics, Inventiones Mathematicae, 73,437-443 3(1983).

[24] J. Gauntlett, D. Martelli, J. Sparks, S-T. Yau: Obstructions to the existence of Sasaki-Einstein metrics. Comm. Math. Phys. 273 (2007), no. 3, 803-827.

[25] Alessandro Ghigi; János Kollár, Kähler-Einstein metrics on orbifolds and Einstein metrics on spheres. Comment. Math. Helv. 82 (2007), no. 4, 877-902.

[26] P. Griffiths, J. Harris: Principles of algebraic geometry, 1978.

[27] B.Guan: The Dirichlet problem for complex Monge-Ampère equations and regularity of the pluri-complex Green function, Comm. Anal. Geom. 6 (1998), 687-703.
[28] Sławomir Kołodziej: The complex Monge-Ampère equation, Acta Math. 180 (1998), no. 1, 69-117
[29] Robin Hartshorne: Algebraic Geometry. Graduate Texts in Mathematics, No. **52**. Springer-Verlag, New York-Heidelberg, 1977.
[30] Thalia D. Jeffres: Uniqueness of Kähler-Einstein cone metrics, Publ. Math. 44(2000), 437-448.
[31] Thalia D. Jeffres, Rafe Mazzeo, Yanir A. Rubinstein: Kähler-Einstein metrics with edge singularities, with an appendix by Chi Li and Yanir A. Rubinstein. arXiv:1105.5216.
[32] Paul Hacking, Yuri Prokhorov: Smoothable del Pezzo surfaces with quotient singularities. Compositio Mathematica (2010), 146, pp: 169-192.
[33] Chi Li: Greatest lower bounds on the Ricci curvature of toric Fano manifolds, Advances in Mathematics 226 (2011) 4921-2932
[34] Chi Li: On the limit behavior of metrics in continuity method to Kahler-Einstein problem in toric Fano case, arXiv:1012.5229. Accepted by Compositio Mathematica.
[35] Chi Li: Remarks on logarithmic K-stability. arXiv:1104.0428
[36] Chi Li: PhD thesis, 2012. Available at www.math.princeton.edu/ chil.
[37] Chi Li, Chenyang Xu: Special test configurations and K-stability of Fano varieties. arXiv:1111.5398
[38] F. Luo, G. Tian: Liouville equation and spherical convex polytopes, Proc. Amer. Math. Soc. 116(1992), 1119-1129.
[39] Toshiki Mabuchi: K-energy maps integrating Futaki invariants. Tohoku Math. J. (2) Vol 38, No. 4 (1986), 575-593.
[40] R.C. McOwen: Point singularties and conformal metrics on Riemann surfaces. Proc. Amer. Math. Soc. 103(1988), 222-224.
[41] Yuji Odaka, Song Sun: Testing log-K-stability by blowing up formalism. arXiv:1112.1353.
[42] Sean T. Paul: Geometric analysis of Chow Mumford stability. *Advances in Mathematics* 182 (2004): 333-356.
[43] Sean T. Paul: Hyperdiscriminant polytopes, Chow polytopes, and Mabuchi energy asymptotics, Annals of Mathematics, 175(2012), 255-296.
[44] Sean T. Paul, Gang Tian: CM stability and the generalized Futaki invariant I. arXiv: 0605278.
[45] Sean T. Paul, Gang Tian: CM stability and the generalized Futaki invariant II. Astérisque No. **328**, 339-354.
[46] D.H.Phong, and Jacob Sturm: The Dirichlet problem for degenerate complex Monge-Ampere equations. arXiv:0904.1898.
[47] D.H.Phong, Jian Song, Jacob Sturm, and Ben Weinkove: The Moser-Trudinger inequality on Kähler-Einstein manifolds, Americal Journal of Mathematics, Volume 130, Number 4, 2008, 1067-1085.
[48] Julius Ross, and Richard Thomas: A study of the Hilbert-Mumford criterion for the stability of projective varieties. J. Algebraic Geom. 16 (2007), 201-255.
[49] Julius Ross, Richard Thomas: Weighted projective embeddings, stability of orbifolds and constant scalar curvature Kähler metrics, Jour. Diff. Geom. 88, 109-160, 2011.
[50] Yalong Shi, and Xiaohua Zhu: Kähler-Ricci solitons on toric Fano orbifolds, arXiv:1102.2764.
[51] Jian Song, Xiaowei Wang: The greatest Ricci lower bound, conical Einstein metrics and the Chern number inequality, in preprint.
[52] Song Sun: Note on K-stability of pairs. arXiv:1108.4603, To appear in Math. Annalen.
[53] Song Sun, Yuanqi Wang: On the Kähler-Ricci flow near a Kähler-Einstein metric, arxiv: 1004.2018.
[54] Szekelyhidi, G.: Greatest lower bounds on the Ricci curvature of Fano manifolds, Compositio Math. 147 (2011), 319-331.
[55] Gang Tian: On Kähler-Einstein metrics on certain Kähler manifolds with $c_1(M) > 0$, Invent. Math., 89 (1987) 225-246
[56] Gang Tian: On stability of the tangent bundles of Fano varieties, Internat. J. Math. 3, 3(1992), 401-413
[57] Gang Tian: Kähler-Einstein metrics on algebraic manifolds, in: Transcendental methods in algebraic geometry (Cetraro 1994), Lecture Notes in Math. 1646, pp. 143-185.
[58] Gang Tian: The K-energy on hypersurfaces and stability. *Communications in Analysis and Geometry* 2, no. 2 (1994): 239-265.
[59] Gang Tian: Kähler-Einstein metrics on algebraic manifolds. Lecture notes in Mathematics, 1996, Volume 1646/1996, 143-185.
[60] Gang Tian: Kähler-Einstein metrics with positive scalar curvature. Invent. math. 137, 1-37 (1997)
[61] Gang Tian: Canonical Metrics on Kähler Manifolds, Birkhauser, 1999
[62] Gang Tian: Existence of Einstein metrics on Fano manifolds. Progress in Mathematics, 2012, Volume 297, Part 1, 119-159.
[63] M. Troyanov: Prescribing curvature on compact surfaces with conic singularities, Trans. Amer. Math. Soc. 324 (1991), 793-821.
[64] Xujia Wang, Xiaohua Zhu: Kähler-Ricci solitons on toric manifolds with positive first Chern class. Advances in Math. 188 (2004) 87-103

[65] Shing-Tung Yau: On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I, Comm. Pure Appl. Math. 31 (1978) 339-441.

[66] Shouwu Zhang: Heights and reductions of Semi-stable varieties. *Compositio Mathematica* 104 (1996): 77-105.

Chi Li
Department of Mathematics, Princeton University, Princeton, NJ 08544, USA.
E-mail: chil@math.princeton.edu

Song Sun
Department of Mathematics, Imperial College, London SW7 2AZ, U.K.
E-mail: s.sun@imperial.ac.uk